Case No. 1:15-cv-02362-RBJ Document 64-25 filed 09/11/17 USDC Colorado pg 1 of 26

overview in 2003, *J Pediatr Surg* 2004;**39**:144–5.
62 **Warne GL**, Grover S, Zajac JD. Hormonal therapies for individuals with intersex conditions: protocol for use. *Treat Endocrinol* 2005;**4**:19–29.
63 **Rogol AD**. New facets of androgen replacement therapy during childhood and adolescence. *Expert Opin Pharmacother* 2005;**6**:1319–36.
64 **Ahmed SF**, Tucker P, Mayo A, *et al*. Randomized, crossover comparison study of the short-term effect of oral testosterone undecanoate and intramuscular testosterone depot on linear growth and serum bone alkaline phosphatase. *J Pediatr Endocrinol Metab* 2004;**17**:941–50.
65 **Mayo A**, Macintyre H, Wallace AM, *et al*. Transdermal testosterone application: pharmacokinetics and effects on pubertal status, short-term growth, and bone turnover. *J Clin Endocrinol Metab* 2004;**89**:681–7.
66 **Weidemann W**, Peters B, Romalo G, *et al*. Response to androgen treatment in a patient with partial androgen insensitivity and a mutation in the deoxyribonucleic acid binding domain of the androgen receptor. *J Clin Endocrinol Metab* 2000;**83**:1173–81.
67 **Kazak AE**, Cant MC, Jensen MM, *et al*. Identifying psychosocial risk indicative of subsequent resource use in families of newly diagnosed pediatric oncology patients. *J Clin Oncol* 2003;**21**:3220–5.
68 **Martin CL**, Ruble DN, Szkrybalo J. Cognitive theories of early gender development. *Psychol Bull* 2002;**128**:903–33.
69 **Zucker KJ**. Measurement of psychosexual differentiation. *Arch Sex Behav* 2005;**34**:375–88.
70 **Carmichael P**, Ransley P. Telling children about a physical intersex condition. *Dialogues Pediatr Urol* 2002;**25**:7–8.
71 **Committee on Paediatric AIDS**, American Academy of Pediatrics. Disclosure of illness status to children and adolescents with HIV infection. *Pediatrics* 1999;**103**:164–6.
72 **Money J**. *Sex errors of the body and related syndromes: a guide to counselling children, adolescents, and their families*, 2nd edition. Baltimore: Paul H Brookes Publishing Co, 1994.
73 **Basson R**, Leiblum S, Brotto L, *et al*. Definitions of women's sexual dysfunction reconsidered: advocating expansion and revision. *J Psychosom Obstet Gynecol* 2003;**24**:221–9.
74 **Creighton S**, Alderson J, Brown S, Minto CL. Medical photography: ethics, consent and the intersex patient. *BJU Int* 2002;**89**:67–71.
75 **Ursano RJ**, Bell C, Eth S, *et al*. Work Group on ASD and PTSD; Steering Committee on Practice Guidelines. Practice guidelines for the treatment of patients with acute stress disorder and posttraumatic stress disorder. *Am J Psychiatry* 2004;**161**(11 suppl):3–31.
76 **Kuhnle U**, Bullinger M Outcome of congenital adrenal hyperplasia. *Pediatr Surg Int* 1997;**12**:511–15.
77 **Warne G**, Grover S, Hutson J, *et al*. Murdoch Childrens Research Institute Sex Study Group. A long-term outcome study of intersex conditions. *J Pediatr Endocrinol Metab* 2005;**18**:555–67.
78 **Creighton SM**. Long-term outcome of feminization surgery: the London experience. *BJU Int* 2004;**93**(suppl 3):44–6.
79 **Steiner E**, Woernie F. Carcinoma of the neovagina: case report and review of the literature. *Gynecol Oncol* 2002;**84**:171–5.
80 **Schober JM**. Long-term outcomes of feminizing genitoplasty for intersex. In: *Pediatric surgery and urology: long-term outcomes*. London: WB Saunders Co (in press)).
81 **Wisniewski AB**, Migeon CJ, Meyer-Bahlburg HF, *et al*. Complete androgen insensitivity syndrome: long-term medical, surgical, and psychosexual outcome. *J Clin Endocrinol Metab* 2000;**85**:2664–9.
82 **Honecker F**, Stoop H, de Krijger RR, *et al*. Pathobiological implications of the expression of markers of testicular carcinoma in situ by fetal germ cells. *J Pathol* 2004;**203**:849–57.
83 **Cools M**, Van Aerde K, Kersemaekers AM, *et al*. Morphological and immunohistochemical differences between gonadal maturation delay and early germ cell neoplasia in patients with undervirilization syndromes. *J Clin Endocrinol Metab* 2005;**90**:5295–303.
84 **Cools M**, Honecker F, Stoop H, *et al*. Maturation delay of germ cells in trisomy 21 fetuses results in increased risk for the development of testicular germ cell tumors. *Hum Pathol* 2006;**37**:101–11.
85 **Ramani P**, Yeung CK, Habeebu SS. Testicular intratubular germ cell neoplasia in children and adults with intersex. *Am J Surg Pathol* 1993;**17**:1124–33.
86 **Hannema SE**, Scott IS, Rajperts-De Meyts E, *et al*. Testicular development in the complete androgen insensitivity syndrome. *J Pathol* 2006;**208**:518–27.
87 **Warne GL**, Bhatia V. Intersex, East and West. In:Sytsma S, editor. *Ethics and intersex*. Berlin: Springer, 2006:183–205.
88 **Achermann JC**, Ozisik G, Meeks JJ, *et al*. Genetic causes of human reproductive disease. *J Clin Endocrinol Metab* 2002;**87**:2447–54.
89 **Small CL**, Shima JE, Uzumcu M, *et al*. Profiling gene expression during the differentiation and development of the murine embryonic gonad. *Biol Reprod* 2005;**72**:492–501.
90 **Martin PL**. Moving toward an international standard in informed consent: the impact of intersexuality and the internet on the standard of care. *Duke J Gend Law Policy* 2002;**9**:135–69.
91 **Department of Health**. *Reference guide to consent for examination or treatment*. London, DoH, 2001 (www.doh.gov.uk/consent).
92 **Sentencia SU-337/99**, May 12, 1999; T-551/99, Aug 2 1999.

APPENDIX 1

ROLE OF SUPPORT GROUPS

The value of peer and parent support for many chronic medical conditions is widely accepted, and DSDs, being lifelong conditions which affect developmental tasks at many stages of life, are no exception.

Those affected by DSDs and parent members value the following:

- Peer support ends isolation and stigma, providing a context in which conditions are put into perspective, and where intimate issues of concern can be discussed safely with someone who has "been there."
- Children who form relationships with peers and affected adults early in their lives benefit from a feeling of normalcy early on, with support in place well before adolescence. Adolescents often resist attempts to introduce them to peer support.
- Support groups can help families and consumers find the best quality care.

While clinical practice may focus on gender and genital appearance as key outcomes, stigma and experiences associated with having a DSD (both within and outside the medical environment) are more salient issues for many affected people.

Support groups complement the work of the health care team and, together, can help improve services. Initiatives by support groups have led to improvements in management of DSD and research directed towards clinically relevant issues. Dialogue between health care professionals and support groups, and collaboration as partners is to be encouraged.

APPENDIX 2

LEGAL ISSUES

Basic principles of medical law will remain, even as research and clinical experience evolve in aetiology, diagnosis, and treatment. This appendix draws on practice in three countries on standards of medical negligence and patient informed consent. In the USA, the medical profession sets standards of care based on prevailing medical custom.[90] However, a treatment may also be that used by a respected minority of practitioners.

USA

Informed consent in the USA was founded on the principle of battery, whereby it is an offence to violate another person's bodily integrity without consent. Nowadays, most states are concerned with negligent non-disclosure to the patient. The standard of adequate disclosure may be physician based, requiring conduct of a reasonable practitioner, or it may be patient based, asking what a reasonable patient would find material. Physician based disclosure must include information about risks, alternatives, outcomes, and prognosis, with or without treatment.

US courts assume that parents know what is best for their child when parental authority applies to consent for the child (substituted judgement). Parental decisions are deferred to except in situations where potentially life saving treatment is withheld. Consent to treatment by a child is dependent on an understanding of its nature and consequences.

United Kingdom

Medical negligence in the United Kingdom defines treatment that falls below the standard expected of a reasonably competent practitioner. The standard of proof in court is whether negligence is demonstrated on the balance of probabilities. It is incumbent on the practitioner to demonstrate that treatment was consistent with a rationally defensible body of medical opinion. A shift in parental prerogative to consent to treatment was reflected in the Children Act 1989, in which parental rights were replaced by parental responsibilities.[91] UK courts can intervene with orders made requiring or preventing a specific action related to the child. Age is not a barrier to

Zzyym v. Tillerson - AR 0644

Case No. 1:15-cv-02362-RBJ Document 64-25 filed 09/11/17 USDC Colorado pg 2 of 26

informed consent, providing that a minor demonstrates an understanding of the issues sufficient to have the capacity to consent.

Colombia

Colombian law is noted for a reasoned set of guidelines advanced by the highest court in cases of DSD.[92] A protocol was formulated for parental and physician intervention. The process of consent requires "qualified and persistent informed consent" over an extended period of time. Authorisation is given in stages to allow time for the parents to come to terms with their child's condition. The court aimed to strike a balance between parental autonomy for those who did and those who did not want early surgery for their child, until there was clear evidence of harm in deferring surgery until the child was competent to decide. Parents cannot consent for children over 5 years of age, as by then children are deemed to have identified with a gender and so are considered to be autonomous.

IMAGES IN PAEDIATRICS

doi: 10.1136/adc.2006.093682

Intraoral graphite tattoo

Graphite pencils, in addition to their usual role as teaching tools, may cause traumatic injury and foreign body reaction, especially during early childhood.

A 5 year old healthy female patient was referred from the paediatric clinic because of her father's concern about a blue lesion on the upper gum which has been increasing in size over four months. Intraoral examination showed an asymptomatic, firm, well defined, scalloped, blue–black macule measuring 15 mm×5 mm involving the attached gingiva and extending apically to the mucogingival junction in the area labial to the maxillary left primary central and lateral incisors (fig 1). There was no associated inflammation and the lesion failed to respond to the blanching test. A periapical radiograph of the anterior maxillary region did not show any pathological changes. Differential diagnoses of foreign body pigmentation, melanocytic nevi, and malignant melanoma were considered. A full mucoperiosteal flap was raised, revealing abundance of granulation tissue, destruction of labial cortical bone to the central incisor, and residues of solid black granules (fig 2). Histopathology showed mild chronic inflammatory cell infiltrate with multinucleated giant cells. There was no evidence of cellular atypia and solid granules were consistent with pencil graphite.



Figure 1 Intraoral view showing the macule labial to maxillary left primary central and lateral incisors.



Figure 2 Destruction of alveolar labial bone and presence of black granules.

The most common causes of exogenous localised oral pigmentation are amalgam tattoo, followed by graphite.[1 2] Intraoral graphite implantation is common in the anterior palates of younger children; biopsy is mandatory to rule out malignancy.[2]

F B Rihani
Prince Rashid Bin AL-Hassan Hospital, Royal Medical Services, Jordan
D M Da'ameh
Queen Alia Hospital, Royal Medical Services, Jordan

Correspondence to: Dr F B Rihani, Irbid, AL-Huson, PO Box 434, Irbid 21510, Jordan; fbrihani@yahoo.com

Competing interests: none

Consent was obtained for publication of the figures

References

1 **Witman PM**, Rogers RS 3rd. Pediatric oral medicine. *Dermatol Clin* 2003;**21**:157–70.
2 **Kauzman A**, Pavone M, Blanas N, *et al*. Pigmented lesions of the oral cavity: review, differential diagnosis, and case presentations. *J Can Dent Assoc* 2004;**70**:682–3.

Zzyym v. Tillerson - AR 0645



www.wpath.org


Standards of Care

for the Health of Transsexual, Transgender, and Gender Nonconforming People

---

The World Professional Association for Transgender Health

7th Version



WPATH WORLD PROFESSIONAL ASSOCIATION for TRANSGENDER HEALTH

Standards of Care

for the Health of Transsexual, Transgender, and Gender Nonconforming People

The World Professional Association for Transgender Health

7th Version[1] | www.wpath.org

1 This is the seventh version of the Standards of Care. The original SOC were published in 1979. Previous revisions were in 1980, 1981, 1990, 1998, and 2001.

Table of Contents

I. Purpose and Use of the Standards of Care 1
II. Global Applicability of the Standards of Care 3
III. The Difference between Gender Nonconformity and Gender Dysphoria 4
IV. Epidemiologic Considerations 6
V. Overview of Therapeutic Approaches for Gender Dysphoria 8
VI. Assessment and Treatment of Children and Adolescents with Gender Dysphoria 10
VII. Mental Health 21
VIII. Hormone Therapy 33
IX. Reproductive Health 50
X. Voice and Communication Therapy 52
XI. Surgery 54
XII. Postoperative Care and Follow-Up 64
XIII. Lifelong Preventive and Primary Care 65
XIV. Applicability of the *Standards of Care* to People Living in Institutional Environments . . . 67
XV. Applicability of the *Standards of Care* to People With Disorders of Sex Development . . . 69

References 72

Appendices:

A. Glossary 95
B. Overview of Medical Risks of Hormone Therapy 97
C. Summary of Criteria for Hormone Therapy and Surgeries 104
D. Evidence for Clinical Outcomes of Therapeutic Approaches 107
E. Development Process for the *Standards of Care, Version 7* 109

Purpose and Use of the Standards of Care

The World Professional Association for Transgender Health (WPATH)[1] is an international, multidisciplinary, professional association whose mission is to promote evidence-based care, education, research, advocacy, public policy, and respect for transgender health. The vision of WPATH is to bring together diverse professionals dedicated to developing best practices and supportive policies worldwide that promote health, research, education, respect, dignity, and equality for transsexual, transgender, and gender nonconforming people in all cultural settings.

One of the main functions of WPATH is to promote the highest standards of health care for individuals through the articulation of *Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming People*. The *SOC* are based on the best available science and expert professional consensus.[2] Most of the research and experience in this field comes from a North American and Western European perspective; thus, adaptations of the *SOC* to other parts of the world are necessary. Suggestions for ways of thinking about cultural relativity and cultural competence are included in this version of the *SOC*.

The overall goal of the *SOC* is to provide clinical guidance for health professionals to assist transsexual, transgender, and gender nonconforming people with safe and effective pathways to achieving lasting personal comfort with their gendered selves, in order to maximize their overall health, psychological well-being, and self-fulfillment. This assistance may include primary care, gynecologic and urologic care, reproductive options, voice and communication therapy, mental health services (e.g., assessment, counseling, psychotherapy), and hormonal and surgical treatments. While this is primarily a document for health professionals, the *SOC* may also be used by individuals, their families, and social institutions to understand how they can assist with promoting optimal health for members of this diverse population.

WPATH recognizes that health is dependent upon not only good clinical care but also social and political climates that provide and ensure social tolerance, equality, and the full rights of citizenship. Health is promoted through public policies and legal reforms that promote tolerance and equity

1 Formerly the Harry Benjamin International Gender Dysphoria Association

2 *Standards of Care (SOC), Version 7* represents a significant departure from previous versions. Changes in this version are based upon significant cultural shifts, advances in clinical knowledge, and appreciation of the many health care issues that can arise for transsexual, transgender, and gender nonconforming people beyond hormone therapy and surgery (Coleman, 2009a, b, c, d).

for gender and sexual diversity and that eliminate prejudice, discrimination, and stigma. WPATH is committed to advocacy for these changes in public policies and legal reforms.

The Standards of Care Are Flexible Clinical Guidelines

The *SOC* are intended to be flexible in order to meet the diverse health care needs of transsexual, transgender, and gender nonconforming people. While flexible, they offer standards for promoting optimal health care and guiding the treatment of people experiencing gender dysphoria – broadly defined as discomfort or distress that is caused by a discrepancy between a person's gender identity and that person's sex assigned at birth (and the associated gender role and/or primary and secondary sex characteristics) (Fisk, 1974; Knudson, De Cuypere, & Bockting, 2010b).

As for all previous versions of the *SOC*, the criteria put forth in this document for hormone therapy and surgical treatments for gender dysphoria are clinical guidelines; individual health professionals and programs may modify them. Clinical departures from the *SOC* may come about because of a patient's unique anatomic, social, or psychological situation; an experienced health professional's evolving method of handling a common situation; a research protocol; lack of resources in various parts of the world; or the need for specific harm reduction strategies. These departures should be recognized as such, explained to the patient, and documented through informed consent for quality patient care and legal protection. This documentation is also valuable for the accumulation of new data, which can be retrospectively examined to allow for health care – and the *SOC* – to evolve.

The *SOC* articulate standards of care but also acknowledge the role of making informed choices and the value of harm reduction approaches. In addition, this version of the *SOC* recognizes and validates various expressions of gender that may not necessitate psychological, hormonal, or surgical treatments. Some patients who present for care will have made significant self-directed progress towards gender role changes, transition, or other resolutions regarding their gender identity or gender dysphoria. Other patients will require more intensive services. Health professionals can use the *SOC* to help patients consider the full range of health services open to them, in accordance with their clinical needs and goals for gender expression.

II

Global Applicability of the Standards of Care

While the *SOC* are intended for worldwide use, WPATH acknowledges that much of the recorded clinical experience and knowledge in this area of health care is derived from North American and Western European sources. From place to place, both across and within nations, there are differences in all of the following: social attitudes towards transsexual, transgender, and gender nonconforming people; constructions of gender roles and identities; language used to describe different gender identities; epidemiology of gender dysphoria; access to and cost of treatment; therapies offered; number and type of professionals who provide care; and legal and policy issues related to this area of health care (Winter, 2009).

It is impossible for the *SOC* to reflect all of these differences. In applying these standards to other cultural contexts, health professionals must be sensitive to these differences and adapt the *SOC* according to local realities. For example, in a number of cultures, gender nonconforming people are found in such numbers and living in such ways as to make them highly socially visible (Peletz, 2006). In settings such as these, it is common for people to initiate a change in their gender expression and physical characteristics while in their teens, or even earlier. Many grow up and live in a social, cultural, and even linguistic context quite unlike that of Western cultures. Yet almost all experience prejudice (Peletz, 2006; Winter, 2009). In many cultures, social stigma towards gender nonconformity is widespread and gender roles are highly prescriptive (Winter et al., 2009). Gender nonconforming people in these settings are forced to be hidden, and therefore may lack opportunities for adequate health care (Winter, 2009).

The *SOC* are not intended to limit efforts to provide the best available care to all individuals. Health professionals throughout the world – even in areas with limited resources and training opportunities – can apply the many core principles that undergird the *SOC*. These principles include the following: Exhibit respect for patients with nonconforming gender identities (do not pathologize differences in gender identity or expression); provide care (or refer to knowledgeable colleagues) that affirms patients' gender identities and reduces the distress of gender dysphoria, when present; become knowledgeable about the health care needs of transsexual, transgender, and gender nonconforming people, including the benefits and risks of treatment options for gender dysphoria; match the treatment approach to the specific needs of patients, particularly their goals for gender expression and need for relief from gender dysphoria; facilitate access to appropriate care; seek patients' informed consent before providing treatment; offer continuity of care; and be prepared to support and advocate for patients within their families and communities (schools, workplaces, and other settings).

Terminology is culturally and time-dependent and is rapidly evolving. It is important to use respectful language in different places and times, and among different people. As the *SOC* are translated into other languages, great care must be taken to ensure that the meanings of terms are accurately translated. Terminology in English may not be easily translated into other languages, and vice versa. Some languages do not have equivalent words to describe the various terms within this document; hence, translators should be cognizant of the underlying goals of treatment and articulate culturally applicable guidance for reaching those goals.

III

The Difference Between Gender Nonconformity and Gender Dysphoria

Being Transsexual, Transgender, or Gender Nonconforming Is a Matter of Diversity, Not Pathology

WPATH released a statement in May 2010 urging the de-psychopathologization of gender nonconformity worldwide (WPATH Board of Directors, 2010). This statement noted that "the expression of gender characteristics, including identities, that are not stereotypically associated with one's assigned sex at birth is a common and culturally-diverse human phenomenon [that] should not be judged as inherently pathological or negative."

Unfortunately, there is stigma attached to gender nonconformity in many societies around the world. Such stigma can lead to prejudice and discrimination, resulting in "minority stress" (I. H. Meyer, 2003). Minority stress is unique (additive to general stressors experienced by all people), socially based, and chronic, and may make transsexual, transgender, and gender nonconforming individuals more vulnerable to developing mental health concerns such as anxiety and depression (Institute of Medicine, 2011). In addition to prejudice and discrimination in society at large, stigma can contribute to abuse and neglect in one's relationships with peers and family members, which in turn can lead to psychological distress. However, these symptoms are socially induced and are not inherent to being transsexual, transgender, or gender nonconforming.

Gender Nonconformity Is Not the Same as Gender Dysphoria

Gender nonconformity refers to the extent to which a person's gender identity, role, or expression differs from the cultural norms prescribed for people of a particular sex (Institute of Medicine, 2011). *Gender dysphoria* refers to discomfort or distress that is caused by a discrepancy between a person's gender identity and that person's sex assigned at birth (and the associated gender role and/or primary and secondary sex characteristics) (Fisk, 1974; Knudson, De Cuypere, & Bockting, 2010b). Only *some* gender nonconforming people experience gender dysphoria at *some* point in their lives.

Treatment is available to assist people with such distress to explore their gender identity and find a gender role that is comfortable for them (Bockting & Goldberg, 2006). Treatment is individualized: What helps one person alleviate gender dysphoria might be very different from what helps another person. This process may or may not involve a change in gender expression or body modifications. Medical treatment options include, for example, feminization or masculinization of the body through hormone therapy and/or surgery, which are effective in alleviating gender dysphoria and are medically necessary for many people. Gender identities and expressions are diverse, and hormones and surgery are just two of many options available to assist people with achieving comfort with self and identity.

Gender dysphoria can in large part be alleviated through treatment (Murad et al., 2010). Hence, while transsexual, transgender, and gender nonconforming people may experience gender dysphoria at some point in their lives, many individuals who receive treatment will find a gender role and expression that is comfortable for them, even if these differ from those associated with their sex assigned at birth, or from prevailing gender norms and expectations.

Diagnoses Related to Gender Dysphoria

Some people experience gender dysphoria at such a level that the distress meets criteria for a formal diagnosis that might be classified as a mental disorder. Such a diagnosis is not a license for stigmatization or for the deprivation of civil and human rights. Existing classification systems such as the *Diagnostic Statistical Manual of Mental Disorders (DSM)* (American Psychiatric Association, 2000) and the *International Classification of Diseases* (*ICD*) (World Health Organization, 2007) define hundreds of mental disorders that vary in onset, duration, pathogenesis, functional disability, and treatability. All of these systems attempt to classify clusters of symptoms and conditions, not the individuals themselves. A disorder is a description of something with which a person might struggle, not a description of the person or the person's identity.

Thus, transsexual, transgender, and gender nonconforming individuals are not inherently disordered. Rather, the distress of gender dysphoria, when present, is the concern that might be diagnosable and for which various treatment options are available. The existence of a diagnosis for such dysphoria often facilitates access to health care and can guide further research into effective treatments.

Research is leading to new diagnostic nomenclatures, and terms are changing in both the *DSM* (Cohen-Kettenis & Pfäfflin, 2010; Knudson, De Cuypere, & Bockting, 2010b; Meyer-Bahlburg, 2010; Zucker, 2010) and the *ICD*. For this reason, familiar terms are employed in the *SOC* and definitions are provided for terms that may be emerging. Health professionals should refer to the most current diagnostic criteria and appropriate codes to apply in their practice areas.

IV
Epidemiologic Considerations

Formal epidemiologic studies on the incidence[3] and prevalence[4] of transsexualism specifically or transgender and gender nonconforming identities in general have not been conducted, and efforts to achieve realistic estimates are fraught with enormous difficulties (Institute of Medicine, 2011; Zucker & Lawrence, 2009). Even if epidemiologic studies established that a similar proportion of transsexual, transgender, or gender nonconforming people existed all over the world, it is likely that cultural differences from one country to another would alter both the behavioral expressions of different gender identities and the extent to which gender dysphoria – distinct from one's gender identity – is actually occurring in a population. While in most countries, crossing normative gender boundaries generates moral censure rather than compassion, there are examples in certain cultures of gender nonconforming behaviors (e.g., in spiritual leaders) that are less stigmatized and even revered (Besnier, 1994; Bolin, 1988; Chiñas, 1995; Coleman, Colgan, & Gooren, 1992; Costa & Matzner, 2007; Jackson & Sullivan, 1999; Nanda, 1998; Taywaditep, Coleman, & Dumronggittigule, 1997).

For various reasons, researchers who have studied incidence and prevalence have tended to focus on the most easily counted subgroup of gender nonconforming individuals: transsexual individuals who experience gender dysphoria and who present for gender-transition-related care at specialist gender clinics (Zucker & Lawrence, 2009). Most studies have been conducted in European

3 **incidence**—the number of new cases arising in a given period (e.g., a year)
4 **prevalence**—the number of individuals having a condition, divided by the number of people in the general population

Zzyym v. Tillerson - AR 0655

countries such as Sweden (Wålinder, 1968, 1971), the United Kingdom (Hoenig & Kenna, 1974), the Netherlands (Bakker, Van Kesteren, Gooren, & Bezemer, 1993; Eklund, Gooren, & Bezemer, 1988; van Kesteren, Gooren, & Megens, 1996), Germany (Weitze & Osburg, 1996), and Belgium (De Cuypere et al., 2007). One was conducted in Singapore (Tsoi, 1988).

De Cuypere and colleagues (2007) reviewed such studies, as well as conducted their own. Together, those studies span 39 years. Leaving aside two outlier findings from Pauly in 1968 and Tsoi in 1988, ten studies involving eight countries remain. The prevalence figures reported in these ten studies range from 1:11,900 to 1:45,000 for male-to-female individuals (MtF) and 1:30,400 to 1:200,000 for female-to-male (FtM) individuals. Some scholars have suggested that the prevalence is much higher, depending on the methodology used in the research (for example, Olyslager & Conway, 2007).

Direct comparisons across studies are impossible, as each differed in their data collection methods and in their criteria for documenting a person as transsexual (e.g., whether or not a person had undergone genital reconstruction, versus had initiated hormone therapy, versus had come to the clinic seeking medically-supervised transition services). The trend appears to be towards higher prevalence rates in the more recent studies, possibly indicating increasing numbers of people seeking clinical care. Support for this interpretation comes from research by Reed and colleagues (2009), who reported a doubling of the numbers of people accessing care at gender clinics in the United Kingdom every five or six years. Similarly, Zucker and colleagues (2008) reported a four- to five-fold increase in child and adolescent referrals to their Toronto, Canada clinic over a 30-year period.

The numbers yielded by studies such as these can be considered minimum estimates at best. The published figures are mostly derived from clinics where patients met criteria for severe gender dysphoria and had access to health care at those clinics. These estimates do not take into account that treatments offered in a particular clinic setting might not be perceived as affordable, useful, or acceptable by all self-identified gender dysphoric individuals in a given area. By counting only those people who present at clinics for a specific type of treatment, an unspecified number of gender dysphoric individuals are overlooked.

Other clinical observations (not yet firmly supported by systematic study) support the likelihood of a higher prevalence of gender dysphoria: (i) Previously unrecognized gender dysphoria is occasionally diagnosed when patients are seen with anxiety, depression, conduct disorder, substance abuse, dissociative identity disorders, borderline personality disorder, sexual disorders, and disorders of sex development (Cole, O'Boyle, Emory, & Meyer III, 1997). (ii) Some crossdressers, drag queens/kings or female/male impersonators, and gay and lesbian individuals may be experiencing gender dysphoria (Bullough & Bullough, 1993). (iii) The intensity of some people's gender dysphoria fluctuates below and above a clinical threshold (Docter, 1988). (iv) Gender nonconformity among FtM individuals tends to be relatively invisible in many cultures, particularly to Western health

professionals and researchers who have conducted most of the studies on which the current estimates of prevalence and incidence are based (Winter, 2009).

Overall, the existing data should be considered a starting point, and health care would benefit from more rigorous epidemiologic study in different locations worldwide.

V

Overview of Therapeutic Approaches for Gender Dysphoria

Advancements in the Knowledge and Treatment of Gender Dysphoria

In the second half of the 20th century, awareness of the phenomenon of gender dysphoria increased when health professionals began to provide assistance to alleviate gender dysphoria by supporting changes in primary and secondary sex characteristics through hormone therapy and surgery, along with a change in gender role. Although Harry Benjamin already acknowledged a spectrum of gender nonconformity (Benjamin, 1966), the initial clinical approach largely focused on identifying who was an appropriate candidate for sex reassignment to facilitate a physical change from male to female or female to male as completely as possible (e.g., Green & Fleming, 1990; Hastings, 1974). This approach was extensively evaluated and proved to be highly effective. Satisfaction rates across studies ranged from 87% of MtF patients to 97% of FtM patients (Green & Fleming, 1990), and regrets were extremely rare (1-1.5% of MtF patients and <1% of FtM patients; Pfäfflin, 1993). Indeed, hormone therapy and surgery have been found to be medically necessary to alleviate gender dysphoria in many people (American Medical Association, 2008; Anton, 2009; The World Professional Association for Transgender Health, 2008).

As the field matured, health professionals recognized that while many individuals need both hormone therapy and surgery to alleviate their gender dysphoria, others need only one of these treatment options and some need neither (Bockting & Goldberg, 2006; Bockting, 2008; Lev, 2004). Often with the help of psychotherapy, some individuals integrate their trans- or cross-gender feelings into the gender role they were assigned at birth and do not feel the need to feminize or masculinize their body. For others, changes in gender role and expression are sufficient to alleviate

gender dysphoria. Some patients may need hormones, a possible change in gender role, but not surgery; others may need a change in gender role along with surgery, but not hormones. In other words, treatment for gender dysphoria has become more individualized.

As a generation of transsexual, transgender, and gender nonconforming individuals has come of age – many of whom have benefitted from different therapeutic approaches – they have become more visible as a community and demonstrated considerable diversity in their gender identities, roles, and expressions. Some individuals describe themselves not as gender nonconforming but as unambiguously cross-sexed (i.e., as a member of the other sex; Bockting, 2008). Other individuals affirm their unique gender identity and no longer consider themselves either male or female (Bornstein, 1994; Kimberly, 1997; Stone, 1991; Warren, 1993). Instead, they may describe their gender identity in specific terms such as transgender, bigender, or genderqueer, affirming their unique experience that may transcend a male/female binary understanding of gender (Bockting, 2008; Ekins & King, 2006; Nestle, Wilchins, & Howell, 2002). They may not experience their process of identity affirmation as a "transition," because they never fully embraced the gender role they were assigned at birth or because they actualize their gender identity, role, and expression in a way that does not involve a change from one gender role to another. For example, some youth identifying as genderqueer have always experienced their gender identity and role as such (genderqueer). Greater public visibility and awareness of gender diversity (Feinberg, 1996) has further expanded options for people with gender dysphoria to actualize an identity and find a gender role and expression that is comfortable for them.

Health professionals can assist gender dysphoric individuals with affirming their gender identity, exploring different options for expression of that identity, and making decisions about medical treatment options for alleviating gender dysphoria.

Options for Psychological and Medical Treatment of Gender Dysphoria

For individuals seeking care for gender dysphoria, a variety of therapeutic options can be considered. The number and type of interventions applied and the order in which these take place may differ from person to person (e.g., Bockting, Knudson, & Goldberg, 2006; Bolin, 1994; Rachlin, 1999; Rachlin, Green, & Lombardi, 2008; Rachlin, Hansbury, & Pardo, 2010). Treatments options include the following:

- Changes in gender expression and role (which may involve living part time or full time in another gender role, consistent with one's gender identity);
- Hormone therapy to feminize or masculinize the body;

- Surgery to change primary and/or secondary sex characteristics (e.g., breasts/chest, external and/or internal genitalia, facial features, body contouring);
- Psychotherapy (individual, couple, family, or group) for purposes such as exploring gender identity, role, and expression; addressing the negative impact of gender dysphoria and stigma on mental health; alleviating internalized transphobia; enhancing social and peer support; improving body image; or promoting resilience.

Options for Social Support and Changes in Gender Expression

In addition (or as an alternative) to the psychological and medical treatment options described above, other options can be considered to help alleviate gender dysphoria, for example:

- Offline and online peer support resources, groups, or community organizations that provide avenues for social support and advocacy;
- Offline and online support resources for families and friends;
- Voice and communication therapy to help individuals develop verbal and non-verbal communication skills that facilitate comfort with their gender identity;
- Hair removal through electrolysis, laser treatment, or waxing;
- Breast binding or padding, genital tucking or penile prostheses, padding of hips or buttocks;
- Changes in name and gender marker on identity documents.

VI

Assessment and Treatment of Children and Adolescents with Gender Dysphoria

There are a number of differences in the phenomenology, developmental course, and treatment approaches for gender dysphoria in children, adolescents, and adults. In children and adolescents, a rapid and dramatic developmental process (physical, psychological, and sexual) is involved and

there is greater fluidity and variability in outcomes, particular in prepubertal children. Accordingly, this section of the *SOC* offers specific clinical guidelines for the assessment and treatment of gender dysphoric children and adolescents.

Differences between Children and Adolescents with Gender Dysphoria

An important difference between gender dysphoric children and adolescents is in the proportion for whom dysphoria persists into adulthood. Gender dysphoria during childhood does not inevitably continue into adulthood.[5] Rather, in follow-up studies of prepubertal children (mainly boys) who were referred to clinics for assessment of gender dysphoria, the dysphoria persisted into adulthood for only 6-23% of children (Cohen-Kettenis, 2001; Zucker & Bradley, 1995). Boys in these studies were more likely to identify as gay in adulthood than as transgender (Green, 1987; Money & Russo, 1979; Zucker & Bradley, 1995; Zuger, 1984). Newer studies, also including girls, showed a 12-27% persistence rate of gender dysphoria into adulthood (Drummond, Bradley, Peterson-Badali, & Zucker, 2008; Wallien & Cohen-Kettenis, 2008).

In contrast, the persistence of gender dysphoria into adulthood appears to be much higher for adolescents. No formal prospective studies exist. However, in a follow-up study of 70 adolescents who were diagnosed with gender dysphoria and given puberty suppressing hormones, all continued with the actual sex reassignment, beginning with feminizing/masculinizing hormone therapy (de Vries, Steensma, Doreleijers, & Cohen-Kettenis, 2010).

Another difference between gender dysphoric children and adolescents is in the sex ratios for each age group. In clinically referred, gender dysphoric children under age 12, the male/female ratio ranges from 6:1 to 3:1 (Zucker, 2004). In clinically referred, gender dysphoric adolescents older than age 12, the male/female ratio is close to 1:1 (Cohen-Kettenis & Pfäfflin, 2003).

As discussed in section IV and by Zucker and Lawrence (2009), formal epidemiologic studies on gender dysphoria – in children, adolescents, and adults – are lacking. Additional research is needed to refine estimates of its prevalence and persistence in different populations worldwide.

5 Gender nonconforming behaviors in children may continue into adulthood, but such behaviors are not necessarily indicative of gender dysphoria and a need for treatment. As described in section III, gender dysphoria is not synonymous with diversity in gender expression.

Phenomenology in Children

Children as young as age two may show features that could indicate gender dysphoria. They may express a wish to be of the other sex and be unhappy about their physical sex characteristics and functions. In addition, they may prefer clothes, toys, and games that are commonly associated with the other sex and prefer playing with other-sex peers. There appears to be heterogeneity in these features: Some children demonstrate extremely gender nonconforming behavior and wishes, accompanied by persistent and severe discomfort with their primary sex characteristics. In other children, these characteristics are less intense or only partially present (Cohen-Kettenis et al., 2006; Knudson, De Cuypere, & Bockting, 2010a).

It is relatively common for gender dysphoric children to have co-existing internalizing disorders such as anxiety and depression (Cohen-Kettenis, Owen, Kaijser, Bradley, & Zucker, 2003; Wallien, Swaab, & Cohen-Kettenis, 2007; Zucker, Owen, Bradley, & Ameeriar, 2002). The prevalence of autistic spectrum disorders seems to be higher in clinically referred, gender dysphoric children than in the general population (de Vries, Noens, Cohen-Kettenis, van Berckelaer-Onnes, & Doreleijers, 2010).

Phenomenology in Adolescents

In most children, gender dysphoria will disappear before or early in puberty. However, in some children these feelings will intensify and body aversion will develop or increase as they become adolescents and their secondary sex characteristics develop (Cohen-Kettenis, 2001; Cohen-Kettenis & Pfäfflin, 2003; Drummond et al., 2008; Wallien & Cohen-Kettenis, 2008; Zucker & Bradley, 1995). Data from one study suggest that more extreme gender nonconformity in childhood is associated with persistence of gender dysphoria into late adolescence and early adulthood (Wallien & Cohen-Kettenis, 2008). Yet many adolescents and adults presenting with gender dysphoria do not report a history of childhood gender nonconforming behaviors (Docter, 1988; Landén, Wålinder, & Lundström, 1998). Therefore, it may come as a surprise to others (parents, other family members, friends, and community members) when a youth's gender dysphoria first becomes evident in adolescence.

Adolescents who experience their primary and/or secondary sex characteristics and their sex assigned at birth as inconsistent with their gender identity may be intensely distressed about it. Many, but not all, gender dysphoric adolescents have a strong wish for hormones and surgery. Increasing numbers of adolescents have already started living in their desired gender role upon entering high school (Cohen-Kettenis & Pfäfflin, 2003).

Among adolescents who are referred to gender identity clinics, the number considered eligible for early medical treatment – starting with GnRH analogues to suppress puberty in the first Tanner stages – differs among countries and centers. Not all clinics offer puberty suppression. If such treatment is offered, the pubertal stage at which adolescents are allowed to start varies from Tanner stage 2 to stage 4 (Delemarre-van de Waal & Cohen-Kettenis, 2006; Zucker et al., in press). The percentages of treated adolescents are likely influenced by the organization of health care, insurance aspects, cultural differences, opinions of health professionals, and diagnostic procedures offered in different settings.

Inexperienced clinicians may mistake indications of gender dysphoria for delusions. Phenomenologically, there is a qualitative difference between the presentation of gender dysphoria and the presentation of delusions or other psychotic symptoms. The vast majority of children and adolescents with gender dysphoria are not suffering from underlying severe psychiatric illness such as psychotic disorders (Steensma, Biemond, de Boer, & Cohen-Kettenis, published online ahead of print January 7, 2011).

It is more common for adolescents with gender dysphoria to have co-existing internalizing disorders such as anxiety and depression, and/or externalizing disorders such as oppositional defiant disorder (de Vries et al., 2010). As in children, there seems to be a higher prevalence of autistic spectrum disorders in clinically referred, gender dysphoric adolescents than in the general adolescent population (de Vries et al., 2010).

Competency of Mental Health Professionals Working with Children or Adolescents with Gender Dysphoria

The following are recommended minimum credentials for mental health professionals who assess, refer, and offer therapy to children and adolescents presenting with gender dysphoria:

1. Meet the competency requirements for mental health professionals working with adults, as outlined in section VII;

2. Trained in childhood and adolescent developmental psychopathology;

3. Competent in diagnosing and treating the ordinary problems of children and adolescents.

Roles of Mental Health Professionals Working with Children and Adolescents with Gender Dysphoria

The roles of mental health professionals working with gender dysphoric children and adolescents may include the following:

1. Directly assess gender dysphoria in children and adolescents (see general guidelines for assessment, below).

2. Provide family counseling and supportive psychotherapy to assist children and adolescents with exploring their gender identity, alleviating distress related to their gender dysphoria, and ameliorating any other psychosocial difficulties.

3. Assess and treat any co-existing mental health concerns of children or adolescents (or refer to another mental health professional for treatment). Such concerns should be addressed as part of the overall treatment plan.

4. Refer adolescents for additional physical interventions (such as puberty suppressing hormones) to alleviate gender dysphoria. The referral should include documentation of an assessment of gender dysphoria and mental health, the adolescent's eligibility for physical interventions (outlined below), the mental health professional's relevant expertise, and any other information pertinent to the youth's health and referral for specific treatments.

5. Educate and advocate on behalf of gender dysphoric children, adolescents, and their families in their community (e.g., day care centers, schools, camps, other organizations). This is particularly important in light of evidence that children and adolescents who do not conform to socially prescribed gender norms may experience harassment in school (Grossman, D'Augelli, & Salter, 2006; Grossman, D'Augelli, Howell, & Hubbard, 2006; Sausa, 2005), putting them at risk for social isolation, depression, and other negative sequelae (Nuttbrock et al., 2010).

6. Provide children, youth, and their families with information and referral for peer support, such as support groups for parents of gender nonconforming and transgender children (Gold & MacNish, 2011; Pleak, 1999; Rosenberg, 2002).

Assessment and psychosocial interventions for children and adolescents are often provided within a multi-disciplinary gender identity specialty service. If such a multidisciplinary service is not available, a mental health professional should provide consultation and liaison arrangements with a pediatric endocrinologist for the purpose of assessment, education, and involvement in any decisions about physical interventions.

Psychological Assessment of Children and Adolescents

When assessing children and adolescents who present with gender dysphoria, mental health professionals should broadly conform to the following guidelines:

1. Mental health professionals should not dismiss or express a negative attitude towards nonconforming gender identities or indications of gender dysphoria. Rather, they should acknowledge the presenting concerns of children, adolescents, and their families; offer a thorough assessment for gender dysphoria and any co-existing mental health concerns; and educate clients and their families about therapeutic options, if needed. Acceptance and removal of secrecy can bring considerable relief to gender dysphoric children/adolescents and their families.

2. Assessment of gender dysphoria and mental health should explore the nature and characteristics of a child's or adolescent's gender identity. A psychodiagnostic and psychiatric assessment – covering the areas of emotional functioning, peer and other social relationships, and intellectual functioning/school achievement – should be performed. Assessment should include an evaluation of the strengths and weaknesses of family functioning. Emotional and behavioral problems are relatively common, and unresolved issues in a child's or youth's environment may be present (de Vries, Doreleijers, Steensma, & Cohen-Kettenis, 2011; Di Ceglie & Thümmel, 2006; Wallien et al., 2007).

3. For adolescents, the assessment phase should also be used to inform youth and their families about the possibilities and limitations of different treatments. This is necessary for informed consent, but also important for assessment. The way that adolescents respond to information about the reality of sex reassignment can be diagnostically informative. Correct information may alter a youth's desire for certain treatment, if the desire was based on unrealistic expectations of its possibilities.

Psychological and Social Interventions for Children and Adolescents

When supporting and treating children and adolescents with gender dysphoria, health professionals should broadly conform to the following guidelines:

1. Mental health professionals should help families to have an accepting and nurturing response to the concerns of their gender dysphoric child or adolescent. Families play an important role in the psychological health and well-being of youth (Brill & Pepper, 2008; Lev, 2004). This also applies to peers and mentors from the community, who can be another source of social support.

2. Psychotherapy should focus on reducing a child's or adolescent's distress related to the gender dysphoria and on ameliorating any other psychosocial difficulties. For youth pursuing sex reassignment, psychotherapy may focus on supporting them before, during, and after reassignment. Formal evaluations of different psychotherapeutic approaches for this situation have not been published, but several counseling methods have been described (Cohen-Kettenis, 2006; de Vries, Cohen-Kettenis, & Delemarre-van de Waal, 2006; Di Ceglie & Thümmel, 2006; Hill, Menvielle, Sica, & Johnson, 2010; Malpas, in press; Menvielle & Tuerk, 2002; Rosenberg, 2002; Vanderburgh, 2009; Zucker, 2006).

Treatment aimed at trying to change a person's gender identity and expression to become more congruent with sex assigned at birth has been attempted in the past without success (Gelder & Marks, 1969; Greenson, 1964), particularly in the long term (Cohen-Kettenis & Kuiper, 1984; Pauly, 1965). Such treatment is no longer considered ethical.

1. Families should be supported in managing uncertainty and anxiety about their child's or adolescent's psychosexual outcomes and in helping youth to develop a positive self-concept.

2. Mental health professionals should not impose a binary view of gender. They should give ample room for clients to explore different options for gender expression. Hormonal or surgical interventions are appropriate for some adolescents, but not for others.

3. Clients and their families should be supported in making difficult decisions regarding the extent to which clients are allowed to express a gender role that is consistent with their gender identity, as well as the timing of changes in gender role and possible social transition. For example, a client might attend school while undergoing social transition only partly (e.g., by wearing clothing and having a hairstyle that reflects gender identity) or completely (e.g., by also using a name and pronouns congruent with gender identity). Difficult issues include whether and when to inform other people of the client's situation, and how others in their lives should respond.

4. Health professionals should support clients and their families as educators and advocates in their interactions with community members and authorities such as teachers, school boards, and courts.

5. Mental health professionals should strive to maintain a therapeutic relationship with gender nonconforming children/adolescents and their families throughout any subsequent social changes or physical interventions. This ensures that decisions about gender expression and the treatment of gender dysphoria are thoughtfully and recurrently considered. The same reasoning applies if a child or adolescent has already socially changed gender role prior to being seen by a mental health professional.

Social Transition in Early Childhood

Some children state that they want to make a social transition to a different gender role long before puberty. For some children, this may reflect an expression of their gender identity. For others, this could be motivated by other forces. Families vary in the extent to which they allow their young children to make a social transition to another gender role. Social transitions in early childhood do occur within some families with early success. This is a controversial issue, and divergent views are held by health professionals. The current evidence base is insufficient to predict the long-term outcomes of completing a gender role transition during early childhood. Outcomes research with children who completed early social transitions would greatly inform future clinical recommendations.

Mental health professionals can help families to make decisions regarding the timing and process of any gender role changes for their young children. They should provide information and help parents to weigh the potential benefits and challenges of particular choices. Relevant in this respect are the previously described relatively low persistence rates of childhood gender dysphoria (Drummond et al., 2008; Wallien & Cohen-Kettenis, 2008). A change back to the original gender role can be highly distressing and even result in postponement of this second social transition on the child's part (Steensma & Cohen-Kettenis, 2011). For reasons such as these, parents may want to present this role change as an exploration of living in another gender role, rather than an irreversible situation. Mental health professionals can assist parents in identifying potential in-between solutions or compromises (e.g., only when on vacation). It is also important that parents explicitly let the child know that there is a way back.

Regardless of a family's decisions regarding transition (timing, extent), professionals should counsel and support them as they work through the options and implications. If parents do not allow their young child to make a gender role transition, they may need counseling to assist them with meeting their child's needs in a sensitive and nurturing way, ensuring that the child has ample possibilities to explore gender feelings and behavior in a safe environment. If parents do allow their young child to make a gender role transition, they may need counseling to facilitate a positive experience for their child. For example, they may need support in using correct pronouns, maintaining a safe and supportive environment for their transitioning child (e.g., in school, peer group settings), and communicating with other people in their child's life. In either case, as a child nears puberty, further assessment may be needed as options for physical interventions become relevant.

Physical Interventions for Adolescents

Before any physical interventions are considered for adolescents, extensive exploration of psychological, family, and social issues should be undertaken, as outlined above. The duration of this exploration may vary considerably depending on the complexity of the situation.

Physical interventions should be addressed in the context of adolescent development. Some identity beliefs in adolescents may become firmly held and strongly expressed, giving a false impression of irreversibility. An adolescent's shift towards gender conformity can occur primarily to please the parents and may not persist or reflect a permanent change in gender dysphoria (Hembree et al., 2009; Steensma et al., published online ahead of print January 7, 2011).

Physical interventions for adolescents fall into three categories or stages (Hembree et al., 2009):

1. *Fully reversible interventions.* These involve the use of GnRH analogues to suppress estrogen or testosterone production and consequently delay the physical changes of puberty. Alternative treatment options include progestins (most commonly medroxyprogesterone) or other medications (such as spironolactone) that decrease the effects of androgens secreted by the testicles of adolescents who are not receiving GnRH analogues. Continuous oral contraceptives (or depot medroxyprogesterone) may be used to suppress menses.

2. *Partially reversible interventions.* These include hormone therapy to masculinize or feminize the body. Some hormone-induced changes may need reconstructive surgery to reverse the effect (e.g., gynaecomastia caused by estrogens), while other changes are not reversible (e.g., deepening of the voice caused by testosterone).

3. *Irreversible interventions.* These are surgical procedures.

A staged process is recommended to keep options open through the first two stages. Moving from one stage to another should not occur until there has been adequate time for adolescents and their parents to assimilate fully the effects of earlier interventions.

Fully Reversible Interventions

Adolescents may be eligible for puberty suppressing hormones as soon as pubertal changes have begun. In order for adolescents and their parents to make an informed decision about pubertal delay, it is recommended that adolescents experience the onset of puberty to at least Tanner Stage 2. Some children may arrive at this stage at very young ages (e.g., 9 years of age). Studies

evaluating this approach only included children who were at least 12 years of age (Cohen-Kettenis, Schagen, Steensma, de Vries, & Delemarre-van de Waal, 2011; de Vries, Steensma et al., 2010; Delemarre-van de Waal, van Weissenbruch, & Cohen Kettenis, 2004; Delemarre-van de Waal & Cohen-Kettenis, 2006).

Two goals justify intervention with puberty suppressing hormones: (i) their use gives adolescents more time to explore their gender nonconformity and other developmental issues; and (ii) their use may facilitate transition by preventing the development of sex characteristics that are difficult or impossible to reverse if adolescents continue on to pursue sex reassignment.

Puberty suppression may continue for a few years, at which time a decision is made to either discontinue all hormone therapy or transition to a feminizing/masculinizing hormone regimen. Pubertal suppression does not inevitably lead to social transition or to sex reassignment.

Criteria for puberty suppressing hormones

In order for adolescents to receive puberty suppressing hormones, the following minimum criteria must be met:

1. The adolescent has demonstrated a long-lasting and intense pattern of gender nonconformity or gender dysphoria (whether suppressed or expressed);

2. Gender dysphoria emerged or worsened with the onset of puberty;

3. Any co-existing psychological, medical, or social problems that could interfere with treatment (e.g., that may compromise treatment adherence) have been addressed, such that the adolescent's situation and functioning are stable enough to start treatment;

4. The adolescent has given informed consent and, particularly when the adolescent has not reached the age of medical consent, the parents or other caretakers or guardians have consented to the treatment and are involved in supporting the adolescent throughout the treatment process.

Regimens, monitoring, and risks for puberty suppression

For puberty suppression, adolescents with male genitalia should be treated with GnRH analogues, which stop luteinizing hormone secretion and therefore testosterone secretion. Alternatively, they may be treated with progestins (such as medroxyprogesterone) or with other medications that block testosterone secretion and/or neutralize testosterone action. Adolescents with female genitalia should be treated with GnRH analogues, which stop the production of estrogens and

progesterone. Alternatively, they may be treated with progestins (such as medroxyprogesterone). Continuous oral contraceptives (or depot medroxyprogesterone) may be used to suppress menses. In both groups of adolescents, use of GnRH analogues is the preferred treatment (Hembree et al., 2009), but their high cost is prohibitive for some patients

During pubertal suppression, an adolescent's physical development should be carefully monitored – preferably by a pediatric endocrinologist – so that any necessary interventions can occur (e.g., to establish an adequate gender appropriate height, to improve iatrogenic low bone marrow density) (Hembree et al., 2009).

Early use of puberty suppressing hormones may avert negative social and emotional consequences of gender dysphoria more effectively than their later use would. Intervention in early adolescence should be managed with pediatric endocrinological advice, when available. Adolescents with male genitalia who start GnRH analogues early in puberty should be informed that this could result in insufficient penile tissue for penile inversion vaginoplasty techniques (alternative techniques, such as the use of a skin graft or colon tissue, are available).

Neither puberty suppression nor allowing puberty to occur is a neutral act. On the one hand, functioning in later life can be compromised by the development of irreversible secondary sex characteristics during puberty and by years spent experiencing intense gender dysphoria. On the other hand, there are concerns about negative physical side effects of GnRH analog use (e.g., on bone development and height). Although the very first results of this approach (as assessed for adolescents followed over 10 years) are promising (Cohen-Kettenis et al., 2011; Delemarre-van de Waal & Cohen-Kettenis, 2006), the long-term effects can only be determined when the earliest treated patients reach the appropriate age.

Partially Reversible Interventions

Adolescents may be eligible to begin feminizing/masculinizing hormone therapy, preferably with parental consent. In many countries, 16-year-olds are legal adults for medical decision-making and do not require parental consent. Ideally, treatment decisions should be made among the adolescent, the family, and the treatment team.

Regimens for hormone therapy in gender dysphoric adolescents differ substantially from those used in adults (Hembree et al., 2009). The hormone regimens for youth are adapted to account for the somatic, emotional, and mental development that occurs throughout adolescence (Hembree et al., 2009).