IN THE UNITED STATES DISTRICT COURT
FOR THE DISTRICT OF DELAWARE

RECKITT BENCKISER PHARMACEUTICALS, INC., RB PHARMACEUTICALS LIMITED, and MONOSOL RX, LLC, Plaintiffs, v. WATSON LABORATORIES, INC., Defendant.	CA. No. 13-cv-1674-RGA **CONSOLIDATED**
RECKITT BENCKISER PHARMACEUTICALS INC., RB PHARMACEUTICALS LIMITED, and MONOSOL RX, LLC, Plaintiffs, v. PAR PHARMACEUTICAL, INC., and INTELGENX TECHNOLOGIES CORP. Defendants.	C.A. No. 1:14-cv-00422-RGA

THIRD AMENDED COMPLAINT FOR PATENT INFRINGEMENT AGAINST PAR PHARMACEUTICAL, INC. AND INTELGENX TECHNOLOGIES CORP.

Plaintiffs Reckitt Benckiser Pharmaceuticals Inc. ("RBP"), RB Pharmaceuticals Limited ("RBP UK"), and MonoSol Rx, LLC ("MonoSol") (collectively, "Plaintiffs") file this Amended Complaint against Defendants Par Pharmaceutical, Inc. ("Par"), and IntelGenX Technologies Corp. ("IGX") and allege as follows:

NATURE OF THE ACTION

1. This is an action for patent infringement arising under the Food and Drug Laws and Patent Laws of the United States, Titles 21 and 35 of the United States Code, respectively,

arising from Defendant Par's submission of an Abbreviated New Drug Application ("ANDA") to the Food and Drug Administration ("FDA") seeking approval to manufacture and sell a generic version of Plaintiff RBP's Suboxone® sublingual film prior to the expiration of United States Patent Nos. 8,475,832 ("the '832 patent") and 8,017,150 ("the '150 patent"), and 8,603,514 ("the '514 patent") (collectively, "the patents-in-suit").

THE PARTIES

2. Plaintiff RBP is a Delaware corporation having a principal place of business at 10710 Midlothian Turnpike, Suite 430, Richmond, Virginia.

3. Plaintiff RBP UK is a United Kingdom corporation having a principal place of business at 103-105 Bath Road, Slough, UK.

4. Plaintiff MonoSol is a Delaware limited liability corporation having a principal place of business at 30 Technology Drive, Warren, New Jersey.

5. On information and belief, Defendant Par is a Delaware corporation having a principal place of business at One Ram Ridge Road, Spring Valley, New York 10977.

6. On information and belief, Defendant IGX is a Delaware corporation having a principal place of business at 6425 Abrams, Ville St-Laurent (Quebec), Canada.

JURISDICTION AND VENUE

7. This Court has subject matter jurisdiction over this action pursuant to 28 U.S.C. §§ 1331 and 1338(a).

8. On information and belief, Par is in the business of making and selling generic pharmaceutical products, which it distributes, markets, and/or sells in Delaware and throughout the United States.

9. Par has previously submitted to the jurisdiction of the United States District Court for the District of Delaware, for example by bringing the patent infringement suit *Par Pharmaceutical Inc. v. Breckenridge Pharmaceutical Inc.*, C.A. No. 13-1114-SLR.

10. This Court has personal jurisdiction over Par because of, *inter alia*, Par's incorporation in Delaware, its continuous and systematic contacts with corporate entities within this judicial district, its previous submission to the jurisdiction of this judicial district, and its marketing and sales activities in this judicial district, including, but not limited to, the substantial, continuous, and systematic distribution, marketing, and/or sales of generic pharmaceutical products to residents of this judicial district.

11. On information and belief, IGX is a drug delivery company focused on the development of oral controlled-release products as well as rapidly disintegrating delivery systems.

12. IGX, directly or through its affiliates, has previously submitted to the jurisdiction of the United States District Court for the District of Delaware, for example by voluntarily substituting in as defendant in the patent infringement suit *Biovail Laboratories International SRL v. IntelGenx Corp.*, C.A. No. 09-605-LPS.

13. This Court has personal jurisdiction over IGX because of, *inter alia*, IGX's incorporation in Delaware, its continuous and systematic contacts with corporate entities within this judicial district, and its previous submission to the jurisdiction of this judicial district.

14. Venue is proper in this district under 28 U.S.C. §§ 1391 and 1400.

THE PATENTS-IN-SUIT

15. Plaintiff RBP UK is the lawful owner of the '832 patent, and Plaintiff RBP is an exclusive licensee of the '832 patent. The '832 patent, entitled "Sublingual and Buccal Film Compositions," duly and legally issued on July 2, 2013, naming Garry L. Myers, Samuel D.

Hillbert, Bill J. Boone, B. Arlie Bogue, Pradeep Sanghvi, and Madhusudan Hariharan as inventors. A true copy of the '832 patent is attached hereto as Exhibit A.

16. Plaintiff MonoSol is the lawful owner of the '150 patent, and Plaintiff RBP is an exclusive licensee of the '150 patent. The '150 patent, entitled "Polyethylene Oxide-Based Films and Drug Delivery Systems Made Therefrom," duly and legally issued on September 13, 2011, naming Robert K. Yang, Richard C. Fuisz, Garry L. Myers, and Joseph M. Fuisz as inventors. A true copy of the '150 patent is attached hereto as Exhibit B.

17. Plaintiff MonoSol is the lawful owner of the '514 patent, and Plaintiff RBP is an exclusive licensee of the '514 patent. The '514 patent, entitled "Uniform Films for Rapid Dissolve Dosage Form Incorporating Taste-Masking Compositions," duly and legally issued on December 10, 2013, naming Robert K. Yang, Richard C. Fuisz, Garry L. Myers, and Joseph M. Fuisz as inventors. A true copy of the '514 patent is attached hereto as Exhibit C.

SUBOXONE® SUBLINGUAL FILM

18. Plaintiff RBP is the holder of New Drug Application ("NDA") No. 22-410 for Suboxone® (buprenorphine hydrochloride and naloxone hydrochloride) sublingual film.

19. On August 30, 2010, the FDA approved NDA No. 22-410 for the manufacture, marketing, and sale of Suboxone® sublingual film for the maintenance treatment of opioid dependence. Plaintiff RBP has sold Suboxone® sublingual film under NDA No. 22-410 since its approval.

20. The patents-in-suit are listed in the FDA's Approved Drug Products with Therapeutic Equivalence Evaluations (the "Orange Book") as covering Suboxone® sublingual film.

THE DRUG APPROVAL PROCESS

21. In 1984, Congress enacted the Drug Price Competition and Patent Term Restoration Act, commonly known as the "Hatch-Waxman Act" and codified at 21 U.S.C. § 355. The Hatch-Waxman Act was intended to balance two important public policy goals. First, Congress wanted to ensure that innovator drug manufacturers would have meaningful patent protection and a period of marketing exclusivity to enable them to recoup their investments in the development of valuable new drugs. Second, Congress sought to ensure that, once the patent protection and marketing exclusivity for these drugs expire, consumers would benefit from the availability of lower priced generic versions of approved drugs.

22. Under 21 U.S.C. § 355(b)(1), the innovator drug manufacturer and NDA applicant is required to submit extensive testing and safety information concerning the drug. In addition, the NDA applicant must submit information on "any patent which claims the drug for which the applicant submitted the application or which claims a method of using such drug and with respect to which a claim of patent infringement could reasonably be asserted." Once the NDA is approved, the FDA lists this patent information in its Approved Drug Products with Therapeutic Equivalence Evaluations, commonly known as the "Orange Book."

23. In contrast, the Hatch-Waxman Act allows ANDA applicants to obtain FDA approval for generic versions of previously-approved drugs without having to repeat the extensive testing required for a new drug application. Under 21 U.S.C. § 355(j), ANDAs can rely on FDA's previous findings of safety and efficacy for an approved drug product, if they demonstrate, among other things, that the generic drug is bioequivalent to the previously-approved drug.

24. When a generic manufacturer submits an ANDA, the FDA conducts a preliminary review of the application to ensure it is sufficiently complete to permit a substantive review. *See* 21 C.F.R. § 314.101(b)(1). "Receipt of an [ANDA] means that FDA has made a threshold determination that the abbreviated application is sufficiently complete to permit a substantive review." *Id.*

25. Under 21 U.S.C. § 355(j)(2)(A)(vii), the ANDA must also include one of the following four certifications with respect to each of the patents listed in the Orange Book for the previously-approved drug product: (i) that the patent information has not been filed ("Paragraph I" certifications); (ii) that the patent has expired ("Paragraph II" certifications); (iii) that the patent will expire on a specific date ("Paragraph III" certifications); or (iv) that the "patent is invalid or will not be infringed by the manufacture, use, or sale of the new drug for which the application is submitted" ("Paragraph IV certifications").

26. Paragraph IV certifications can allow generic manufacturers to obtain FDA approval long before expiration of the patents listed in the Orange Book.

27. If the ANDA includes a Paragraph IV certification, the Hatch-Waxman Act requires the ANDA applicant to give notice ("notice of Paragraph IV certification") to the patent owner of the factual and legal basis for the applicant's opinion that patents listed in the Orange Book are invalid or will not be infringed, "not later than 20 days after the date of the postmark on the notice with which the [FDA] informs the applicant that the application has been filed." 21 U.S.C. § 355(j)(2)(B).

28. The patent owner can file an infringement action within 45 days of receiving the notice of Paragraph IV certification. Such a filing by the patent owner triggers a 30-month injunction or stay of the FDA approval, beginning on the date of receipt of the notice. *See* 21

U.S.C. § 355(j)(5)(B)(iii). This 30-month period is intended to allow time for judicial resolution on the merits of any patent infringement, validity, and/or enforceability claims, before the competitor is allowed entry into the market.

29. The date of receipt of the notice of Paragraph IV certification by the patent owner affects important rights and remedies of the patent owner, including the deadline for filing an infringement action and the start and end dates of the 30-month injunction. By providing a premature notice of Paragraph IV certification to the NDA holder or patent owner, the ANDA applicant attempts to trigger the 30-month injunction earlier and to reach the market sooner than it would otherwise be permitted. In addition, there is a significant risk for the patent owner to miss the 45-day deadline for filing an infringement action, and consequently the opportunity for a 30-month injunction, if the patent owner mistakenly relies on a premature and ineffective notice of Paragraph IV certification.

30. The ANDA applicant may not send the notice of Paragraph IV certification to the patent owner before the FDA "informs the applicant that the application has been filed." 21 U.S.C. § 355(j)(2)(B)(ii)(I); *SB Pharmco Puerto Rico, Inc. v. Mut. Pharm. Co., Inc.*, 552 F. Supp. 2d 500, 508 (E.D. Pa. 2008); *Otsuka Pharm. Co. v. Par Pharm., Inc.*, Docket No. 13-1979 (D. Del. March 10, 2014) (Andrews, J.).

31. Federal regulations also govern the timing of the notice of Paragraph IV certification by directing the generic manufacturer to send such notice "when it receives from FDA an acknowledgment letter stating that its [ANDA] is sufficiently complete to permit a substantive review." 21 C.F.R. § 314.95(b).

DEFENDANTS' ANDA AND NOTICES OF PARAGRAPH IV CERTIFICATION

32. Plaintiffs received a letter from Defendant Par dated July 8, 2013 (the "July 2013 Notice Letter"), stating that ANDA No. 20-5854 contains a certification pursuant to 21 U.S.C. § 355(j)(2)(A)(vii)(IV) (a "Paragraph IV certification") alleging that the '832 and '150 patents are invalid, unenforceable, and/or will not be infringed by the manufacture, use, or sale of the generic product proposed in the ANDA.

33. The July 2013 Notice Letter further states that Defendant Par submitted ANDA No. 20-5854 to the FDA under 21 U.S.C. § 355(j), seeking approval to engage in the commercial manufacture, use, and/or sale of buprenorphine hydrochloride and naloxone hydrochloride sublingual film ("Defendants' generic product") before expiration of the patents-in-suit.

34. On information and belief, ANDA No. 20-5854 refers to and relies on Plaintiff RBP's NDA for Suboxone® sublingual film and purports to contain data showing bioequivalence of Defendants' generic product with Suboxone® sublingual film.

35. On information and belief, ANDA No. 20-5854 was prepared and submitted with the active cooperation, participation, and assistance of, and at least in part for the benefit of, Defendant IGX. On information and belief, if ANDA No. 20-5854 is approved, IGX will actively participate in manufacturing, marketing, and/or selling Defendants' generic product.

36. On information and belief, IGX designed Defendants' generic product that is the subject of Defendant Par's ANDA No. 20-5854.

37. On information and belief, Defendants' generic product that is the subject of Defendant Par's ANDA No. 20-5854 includes IGX's VersaFilm™ drug delivery technology.

38. IGX filed statements with the SEC in 2013 asserting that IGX's "U.S. based co-development and commercialization partner" submitted an ANDA to the FDA for approval of a

generic formulation of Plaintiff RBP's Suboxone® sublingual film, indicated for maintenance treatment of opioid dependence.

39. Plaintiffs filed suit against Defendants in this judicial district on August 20, 2013 (Case No. 1:13-cv-01461-RGA), within 45 days of receiving the July 2013 Notice Letter.

40. Plaintiffs received another letter from Defendant Par dated February 3, 2014 ("the '514 Notice Letter"), stating that ANDA No. 20-5854 contains a Paragraph IV certification pursuant to 21 U.S.C. § 355(j)(2)(A)(vii)(IV), alleging that the '514 patent is invalid and/or will not be infringed by the manufacture, use, or sale of the generic product proposed in the ANDA.

41. Plaintiffs amended their complaint in Case No. 1:13-cv-01461-RGA on February 18, 2014, within 45 days of receiving the '514 Notice Letter, to further assert the infringement of the '514 Patent under 35 U.S.C. § 271(e)(2).

42. On March 4, 2014, Defendants filed an answer to Plaintiffs' first amended complaint in Case No. 1:13-cv-01461-RGA, with counterclaims asserted by Par.

43. Plaintiffs received yet another letter from Defendant Par dated March 25, 2014 (the "March 2014 Notice Letter"), repeating that ANDA No. 20-5854 contains a Paragraph IV certification pursuant to 21 U.S.C. § 355(j)(2)(A)(vii)(IV) alleging that the '832, '150, and '514 patents are invalid, unenforceable, and/or will not be infringed by the manufacture, use, or sale of the generic product proposed in the ANDA.

44. The March 2014 Notice Letter further states that Defendant Par submitted ANDA No. 20-5854 to the FDA under 21 U.S.C. § 355(j), seeking approval to engage in the commercial manufacture, use, and/or sale of Defendants' generic product, i.e. buprenorphine hydrochloride and naloxone hydrochloride sublingual film, before expiration of the patents-in-suit.

45. Unlike the earlier notices of Paragraph IV certifications (which include the July 2013 Notice Letter and the '514 Notice Letter), the March 2014 Notice Letter represents that the FDA "has received [ANDA No. 20-5854] for substantive review."

46. On information and belief, Defendant Par's earlier notices of Paragraph IV certifications, which predate the March 2014 Notice Letter, were sent before receiving the FDA's acknowledgment letter stating that ANDA No. 20-5854 is sufficiently complete to permit a substantive review (the "FDA Acknowledgment Letter").

47. Recent authority from this judicial district demonstrates that Defendant Par's earlier notices of Paragraph IV certifications, which predate the March 2014 Notice Letter, are premature and ineffective to trigger the 45-day period for filing an infringement action and the 30-month injunction on FDA approval of Defendants' ANDA No. 20-5854. *See Otsuka Pharm. Co. v. Par Pharm., Inc.*, Docket No. 13-1979 (D. Del. March 10, 2014) (Andrews, J.).

48. Plaintiffs commenced this action within 45 days of receiving the March 2014 Notice Letter, to preserve their right to a 30-month stay under 21 U.S.C. 355(j)(5)(B)(iii).

49. Plaintiffs received another notice of Paragraph IV certification from Defendant Par, dated April 8, 2014 (the "April 2014 Notice Letter"). The April 2014 Notice Letter states that ANDA No. 20-5854 contains a Paragraph IV certification pursuant to 21 U.S.C. § 355(j)(2)(A)(vii)(IV) alleging that the '832, '150, and '514 patents are invalid, unenforceable, and/or will not be infringed by the manufacture, use, or sale of the generic product proposed in the ANDA.

50. Whereas the March 2014 Notice Letter states that "[t]he active ingredient, strength, and dosage form of the proposed drug product are buprenorphine hydrochloride; naloxone hydrochloride, 8 mg/2 mg (Equivalent Base), and sublingual film," the April 2014

Notice Letter adds the "2 mg/0.5 mg (Equivalent Base)" and "4 mg/1 mg (Equivalent Base)" strengths.

51. Plaintiffs commenced this action within 45 days of receiving the April 2014 Notice Letter, to preserve their right to a 30-month stay under 21 U.S.C. 355(j)(5)(B)(iii) as to the newly added "2 mg/0.5 mg (Equivalent Base)" and "4 mg/1 mg (Equivalent Base)" strengths, which 30-month stay was triggered, at the very earliest, upon Plaintiffs' receipt of the April 2014 Notice Letter.

52. On April 17, 2014, Plaintiffs moved to dismiss their amended complaint and Par's counterclaims in Case No. 1:13-cv-01461-RGA, without prejudice, on the basis that Par admittedly served the Paragraph IV certifications on which that action was based even though its application had not been accepted for filing when it started the litigation process.

53. On May 27, 2014, this Court granted Plaintiffs' motion to dismiss their amended complaint and Par's counterclaims in Case No. 1:13-cv-01461-RGA, without prejudice.

54. Plaintiffs received another notice of Paragraph IV certification from Defendant Par, dated June 13, 2014 (the "June 2014 Notice Letter"). The June 2014 Notice Letter states that ANDA No. 20-5854 contains a Paragraph IV certification pursuant to 21 U.S.C. § 355(j)(2)(A)(vii)(IV) alleging that the '832, '150, and '514 patents are invalid, unenforceable, and/or will not be infringed by the manufacture, use, or sale of the generic product proposed in the ANDA.

55. Whereas the April 2014 Notice Letter states that "[t]he active ingredient, strength, and dosage form of the proposed drug product are buprenorphine hydrochloride; naloxone hydrochloride, 2 mg/0.5 mg (Equivalent Base); 4 mg/1 mg (Equivalent Base); and 8 mg/2 mg

(Equivalent Base), and sublingual film," the June 2014 Notice Letter adds the "12 mg/3 mg (Equivalent Base)" strength.

56. Plaintiffs commenced this action within 45 days of receiving the June 2014 Notice Letter, to preserve their right to a 30-month stay under 21 U.S.C. 355(j)(5)(B)(iii) as to the newly added "12 mg/3 mg (Equivalent Base)" strength, which 30-month stay was triggered, at the very earliest, upon Plaintiffs' receipt of the June 2014 Notice Letter.

COUNT I
(Infringement of the '832 Patent Under 35 U.S.C. § 271(e)(2))

57. Plaintiffs reallege paragraphs 1-56 above as if fully set forth herein.

58. On information and belief, Defendants' generic product is covered by one or more claims of the '832 patent.

59. By filing ANDA No. 20-5854 under 21 U.S.C. § 355(j) for the purposes of obtaining approval to engage in the commercial manufacture, use, sale and/or importation of Defendants' generic product prior to the expiration of the '832 patent, Par has committed an act of infringement of the '832 patent under 35 U.S.C. § 271(e)(2).

60. On information and belief, IGX was actively involved in the preparation and are actively involved in the prosecution before the FDA of ANDA No. 20-5854.

61. IGX's active assistance and involvement with the submission of ANDA No. 20-5854 is an act of infringement of the '832 patent under 35 U.S.C. § 271(e)(2).

62. Plaintiffs are entitled to the relief provided by 35 U.S.C. § 271(e)(4), including, *inter alia*, an order of this Court that the FDA set the effective date of approval for ANDA No. 20-5854 to be a date which is not any earlier than the expiration date of the '832 patent, including any extensions of that date.

COUNT II
(Infringement of the ’150 Patent Under 35 U.S.C. § 271(e)(2))

63. Plaintiffs reallege paragraphs 1-62 above as if fully set forth herein.

64. On information and belief, Defendants’ generic product is covered by one or more claims of the ’150 patent.

65. By filing ANDA No. 20-5854 under 21 U.S.C. § 355(j) for the purposes of obtaining approval to engage in the commercial manufacture, use, sale and/or importation of Defendants’ generic product prior to the expiration of the ’150 patent, Par has committed an act of infringement of the ’150 patent under 35 U.S.C. § 271(e)(2).

66. On information and belief, IGX was actively involved in the preparation and are actively involved in the prosecution before the FDA of ANDA No. 20-5854.

67. IGX’s active assistance and involvement with the submission of ANDA No. 20-5854 is an act of infringement of the ’150 patent under 35 U.S.C. § 271(e)(2).

68. Plaintiffs are entitled to the relief provided by 35 U.S.C. § 271(e)(4), including, inter *alia*, an order of this Court that the FDA set the effective date of approval for ANDA No. 20-5854 to be a date which is not any earlier than the expiration date of the ’150 patent, including any extensions of that date.

COUNT III
(Infringement of the ’514 Patent Under 35 U.S.C. § 271(e)(2))

69. Plaintiffs reallege paragraphs 1-68 above as if fully set forth herein.

70. On information and belief, Defendants’ generic product is covered by one or more claims of the ’514 patent.

71. ANDA No. 20-5854 under 21 U.S.C. § 355(j) seeks to obtain approval to engage in the commercial manufacture, use, sale and/or importation of Defendants’ generic product prior

to the expiration of the ’514 patent. Therefore, Par’s maintenance of this filing constitutes an act of infringement of the ’514 patent under 35 U.S.C. § 271(e)(2).

72. On information and belief, IGX was actively involved in the preparation and are actively involved in the prosecution before the FDA of ANDA No. 20-5854.

73. IGX’s active assistance and involvement with the submission of ANDA No. 20-5854 is an act of infringement of the ’514 patent under 35 U.S.C. § 271(e)(2).

74. Plaintiffs are entitled to the relief provided by 35 U.S.C. § 271(e)(4), including, *inter alia*, an order of this Court that the FDA set the effective date of approval for ANDA No. 20-5854 to be a date which is not any earlier than the expiration date of the ’514 patent, including any extensions of that date.

PRAYER FOR RELIEF

WHEREFORE, Plaintiffs respectfully request that this Court:

A. Enter a judgment that Defendants have infringed each of the patents-in-suit under 35 U.S.C. § 271(e)(2) by submitting and maintaining ANDA No. 20-5854;

B. Enter preliminary and permanent injunctions, restraining and enjoining Defendants, their officers, agents, attorneys, affiliates, divisions, successors and employees, and those acting in privity or concert with them, from engaging in, causing, or inducing the commercial manufacture, use, offer to sell, or sale within the United States, or importation into the United States, of drugs and formulations, or from inducing and/or encouraging the use of methods, claimed in the patents-in-suit;

C. Enter an order that the effective date of any approval of ANDA No. 20-5854 be a date that is not earlier than the expiration of the last to expire of the patents-in-suit, including any extensions thereof and any later expiration of exclusivity associated with those patents;

D. Enter a judgment and order finding that this is an exceptional case within the meaning of 35 U.S.C. § 285 and awarding to Plaintiffs their reasonable attorneys' fees;

E. Enter a judgment granting Plaintiffs compensatory damages in an amount to be determined at trial including both pre-judgment and post-judgment interest if Defendants commercially manufacture, use, offer to sell, or sell in the United States, or import into the United States, Defendants' generic product before the expiration of each patent-in-suit that Defendants are found to infringe, including any extensions; and

F. Order any and all other relief as the Court deems just and proper.

Dated: July 25, 2014

Respectfully submitted,

WOMBLE CARLYLE SANDRIDGE & RICE, LLP

/s/ Mary W. Bourke
Mary W. Bourke (#2356)
Dana K. Severance (#4869)
222 Delaware Avenue, Suite 1501
Wilmington, DE 19801
(302) 252-4320
(302) 252-4330 (Fax)
mbourke@wcsr.com
dseverance@wcsr.com

Attorneys for Plaintiffs

Of Counsel:

Daniel A. Ladow
James M. Bollinger
Timothy P. Heaton
J. Magnus Essunger
TROUTMAN SANDERS LLP
405 Lexington Avenue
New York, NY 10174
(212) 704-6000
(212) 704-6288 (Fax)
Daniel.ladow@troutmansanders.com
James.bollinger@troutmansanders.com
Timothy.heaton@troutmansanders.com
Magnus.Essunger@troutmansanders.com

Troy S. Kleckley
Puja R. Patel
TROUTMAN SANDERS LLP
600 Peachtree Street, NE
Suite 5200
Atlanta, GA 30308
(404) 885-3000

(404) 885-3900 (Fax)
Troy.kleckley@troutmansanders.com
Puja.patel@troutmansanders.com

Attorneys for Plaintiff Reckitt Benckiser Pharmaceuticals, Inc. & RB Pharmaceuticals Limited

James F. Hibey
Timothy C. Bickham
Houda Morad
Stephanie L. Schonewald
STEPTOE & JOHNSON LLP
1330 Connecticut Avenue, NW
Washington DC 20036
(202) 429-3000
(202) 429-3902 (Fax)
jhibey@steptoe.com
tbickham@steptoe.com
hmorad@steptoe.com
sschonew@steptoe.com

Attorneys for Plaintiff MonoSol Rx, LLC

32814212

EXHIBIT A

US008475832B2

(12) **United States Patent**
Myers et al.

(10) **Patent No.: US 8,475,832 B2**
(45) **Date of Patent: Jul. 2, 2013**

(54) **SUBLINGUAL AND BUCCAL FILM COMPOSITIONS**

(75) Inventors: **Garry L. Myers**, Kingsport, TN (US); **Samuel D. Hilbert**, Jonesboro, TN (US); **Bill J. Boone**, Johnson City, TN (US); **B. Arlie Bogue**, New Carlisle, IN (US); **Pradeep Sanghvi**, Schererville, IN (US); **Madhusudan Hariharan**, Munster, IN (US)

(73) Assignee: **RB Pharmaceuticals Limited**, Slough (GB)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 231 days.

(21) Appl. No.: **12/537,571**

(22) Filed: **Aug. 7, 2009**

(65) **Prior Publication Data**

US 2011/0033541 A1 Feb. 10, 2011

(51) **Int. Cl.**

A61F 13/00	(2006.01)
A61K 9/14	(2006.01)
A61K 31/44	(2006.01)

(52) **U.S. Cl.**

USPC **424/435**; 424/422; 424/434; 424/484; 514/282

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

307,537	A	11/1884	Foulks
688,446	A	12/1901	Stempel
2,980,554	A	4/1961	Gentile et al.
3,007,848	A	11/1961	Stroop
3,249,109	A	5/1966	Maeth et al.
3,444,858	A	5/1969	Russell
3,536,809	A	10/1970	Applezweig
3,551,556	A	12/1970	Kliment et al.
3,598,122	A	8/1971	Zaffaroni
3,632,740	A	1/1972	Robinson et al.
3,640,741	A	2/1972	Etes
3,641,237	A	2/1972	Gould et al.
3,731,683	A	5/1973	Zaffaroni
3,753,732	A	8/1973	Boroshok
3,814,095	A	6/1974	Lubens
3,892,905	A	7/1975	Albert
3,911,099	A	10/1975	DeFoney et al.
3,972,995	A	8/1976	Tsuk et al.
3,996,934	A	12/1976	Zaffaroni
4,029,757	A	6/1977	Mlodozeniec et al.
4,029,758	A	6/1977	Mlodozeniec et al.
4,031,200	A	6/1977	Reif
4,123,592	A	10/1978	Rainer et al.
4,128,445	A	12/1978	Sturzenegger et al.
4,136,145	A	1/1979	Fuchs et al.
4,136,162	A	1/1979	Fuchs et al.
4,139,627	A	2/1979	Lane et al.
4,226,848	A	10/1980	Nagai et al.
4,251,400	A	2/1981	Columbus
4,292,299	A	9/1981	Suzuki et al.
4,294,820	A	10/1981	Keith et al.
4,302,465	A	11/1981	Ekenstam et al.
4,307,075	A	12/1981	Martin
4,325,855	A	4/1982	Dickmann et al.
4,373,036	A	2/1983	Chang et al.
4,406,708	A	9/1983	Hesselgren
4,432,975	A	2/1984	Libby
4,438,258	A	3/1984	Graham
4,460,562	A	7/1984	Keith et al.
4,466,973	A	8/1984	Rennie
4,503,070	A	3/1985	Eby, III
4,515,162	A	5/1985	Yamamoto et al.
4,517,173	A	5/1985	Kizawa et al.
4,529,601	A	7/1985	Broberg et al.
4,529,748	A	7/1985	Wienecke
4,562,020	A	12/1985	Hijiya et al.
4,569,837	A	2/1986	Suzuki et al.
4,582,835	A	4/1986	Lewis et al.
4,593,053	A	6/1986	Jevne et al.
4,608,249	A	8/1986	Otsuka et al.
4,615,697	A	10/1986	Robinson
4,623,394	A	11/1986	Nakamura et al.
4,631,837	A	12/1986	Magoon
4,652,060	A	3/1987	Miyake
4,659,714	A	4/1987	Watt-Smith
4,675,009	A	6/1987	Hymes et al.
4,695,465	A	9/1987	Kigasawa et al.
4,704,119	A	11/1987	Shaw et al.
4,713,239	A	12/1987	Babaian et al.
4,713,243	A	12/1987	Schiraldi et al.
4,722,761	A	2/1988	Cartmell et al.
4,740,365	A	4/1988	Yukimatsu et al.
4,748,022	A	5/1988	Busciglio
4,765,983	A	8/1988	Takayanagi et al.
4,772,470	A	9/1988	Inoue et al.
4,777,046	A	10/1988	Iwakura et al.
4,789,667	A	12/1988	Makino et al.
4,849,246	A	7/1989	Schmidt
4,860,754	A	8/1989	Sharik et al.
RE33,093	E	10/1989	Schiraldi et al.
4,876,092	A	10/1989	Mizobuchi et al.
4,876,970	A	10/1989	Bolduc
4,888,354	A	12/1989	Chang et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU	741362	B2	7/1998
DE	2432925	B2	1/1976

(Continued)

OTHER PUBLICATIONS

Lazaridou et al., "Thermophysical proprties of chitosan, chitosan-starch and chitosan-pullulan films near the glass transition," Carbohydrate Polymers 48: 179-190 (2002).

(Continued)

Primary Examiner — Janet Epps-Smith

(74) *Attorney, Agent, or Firm* — Hoffmann & Baron, LLP

(57) **ABSTRACT**

The present invention relates to products and methods for treatment of narcotic dependence in a user. The invention more particularly relates to self-supporting dosage forms which provide an active agent for treating narcotic dependence while providing sufficient buccal adhesion of the dosage form.

19 Claims, No Drawings

U.S. PATENT DOCUMENTS

4,894,232	A	1/1990	Reül et al.
4,900,552	A	2/1990	Sanvordeker et al.
4,900,554	A	2/1990	Yanagibashi et al.
4,900,556	A	2/1990	Wheatley et al.
4,910,247	A	3/1990	Haldar et al.
4,915,950	A	4/1990	Miranda et al.
4,925,670	A	5/1990	Schmidt
4,927,634	A	5/1990	Sorrentino et al.
4,927,636	A	5/1990	Hijiya et al.
4,937,078	A	6/1990	Mezei et al.
4,940,587	A	7/1990	Jenkins et al.
4,948,580	A	8/1990	Browning
4,958,580	A	9/1990	Asaba et al.
4,978,531	A	12/1990	Yamazaki et al.
4,981,693	A	1/1991	Higashi et al.
4,981,875	A	1/1991	Leusner et al.
5,023,082	A	6/1991	Friedman et al.
5,024,701	A	6/1991	Desmarais
5,028,632	A	7/1991	Fuisz
5,045,445	A	9/1991	Schultz
5,047,244	A	9/1991	Sanvordeker et al.
5,064,717	A	11/1991	Suzuki et al.
5,089,307	A	2/1992	Ninomiya et al.
5,118,508	A	6/1992	Kikuchi et al.
5,158,825	A	10/1992	Altwirth
5,166,233	A	11/1992	Kuroya et al.
5,186,938	A	2/1993	Sablotsky et al.
5,229,164	A	7/1993	Pins et al.
5,234,957	A	8/1993	Mantelle
5,271,940	A	12/1993	Cleary et al.
5,272,191	A	12/1993	Ibrahim et al.
5,346,701	A	9/1994	Heiber et al.
5,393,528	A	2/1995	Staab
5,411,945	A	5/1995	Ozaki et al.
5,413,792	A	5/1995	Ninomiya et al.
5,433,960	A	7/1995	Meyers
5,455,043	A	10/1995	Fischel-Ghodsian
5,462,749	A	10/1995	Rencher
5,472,704	A	12/1995	Santus et al.
5,518,902	A	5/1996	Ozaki et al.
5,567,431	A	10/1996	Vert et al.
5,605,696	A	2/1997	Eury et al.
5,620,757	A	4/1997	Ninomiya et al.
5,629,003	A	5/1997	Horstmann et al.
5,681,873	A	10/1997	Norton et al.
5,700,478	A	12/1997	Biegajski et al.
5,700,479	A	12/1997	Lundgren
5,766,332	A	6/1998	Graves et al.
5,766,620	A	6/1998	Heiber et al.
5,766,839	A	6/1998	Johnson et al.
5,800,832	A	9/1998	Tapolsky et al.
5,806,284	A	9/1998	Gifford
5,891,461	A	4/1999	Jona et al.
5,900,247	A	5/1999	Rault et al.
5,948,430	A	9/1999	Zerbe et al.
6,072,100	A	6/2000	Mooney et al.
6,103,266	A	8/2000	Tapolsky et al.
6,153,210	A	11/2000	Roberts et al.
6,159,498	A	12/2000	Tapolsky et al.
6,177,096	B1	1/2001	Zerbe et al.
6,231,957	B1	5/2001	Zerbe et al.
6,264,981	B1	7/2001	Zhang et al.
6,284,264	B1	9/2001	Zerbe et al.
6,375,963	B1	4/2002	Repka et al.
6,503,532	B1	1/2003	Murty et al.
6,667,060	B1	12/2003	Vandecruys et al.
6,800,329	B2	10/2004	Horstmann et al.
6,824,829	B2	11/2004	Berry et al.
7,005,142	B2	2/2006	Leon
7,579,019	B2	8/2009	Tapolsky et al.
2001/0006677	A1	7/2001	McGinity et al.
2001/0022964	A1	9/2001	Leung et al.
2001/0046511	A1	11/2001	Zerbe et al.
2003/0069263	A1	4/2003	Breder et al.
2003/0107149	A1	6/2003	Yang et al.
2003/0124176	A1	7/2003	Hsu et al.
2004/0096569	A1	5/2004	Barkalow et al.
2004/0191302	A1	9/2004	Davidson
2005/0048102	A1	3/2005	Tapolsky et al.
2005/0118217	A1	6/2005	Barnhart et al.
2005/0147658	A1	7/2005	Tapolsky et al.
2005/0192309	A1	9/2005	Palermo et al.
2006/0210610	A1	9/2006	Davidson et al.
2006/0281775	A1	12/2006	Kelly, II et al.
2007/0087036	A1	4/2007	Durschlag et al.
2007/0148097	A1	6/2007	Finn et al.
2008/0254105	A1	10/2008	Tapolsky et al.
2010/0015128	A1	1/2010	Lee et al.
2010/0087470	A1 *	4/2010	Oksche et al. 514/279
2011/0189259	A1	8/2011	Vasisht et al.
2011/0262522	A1	10/2011	Finn et al.

FOREIGN PATENT DOCUMENTS

DE	2449865	B2	4/1976
DE	3630603	C2	3/1988
EP	0219762	B1	12/1990
EP	0259749	B1	8/1991
EP	0200508	B1	10/1991
EP	0241178	B1	1/1992
EP	0273069	B1	10/1992
EP	0250187	B1	9/1993
EP	0452446	B1	12/1993
EP	0381194	B1	8/1994
EP	0440462	B1	12/1994
EP	0514691	B1	3/1996
EP	0598606	B1	6/1999
EP	0949925	B1	10/1999
EP	1110546	A1	6/2001
EP	1897543	A1	3/2008
GB	2447016	A	9/2008
JP	62126950		6/1987
JP	02265444		10/1990
JP	05147140		6/1993
JP	07322812		12/1995
JP	2001279100		10/2001
WO	9105540	A1	5/1991
WO	9215289	A1	9/1992
WO	9505416	A2	2/1995
WO	9518046	A1	7/1995
WO	9817251		4/1998
WO	9955312		11/1999
WO	0018365	A2	4/2000
WO	0042992	A2	7/2000
WO	0170194	A1	9/2001
WO	0191721	A2	12/2001
WO	03/030882	A1	4/2003
WO	03030883	A1	4/2003
WO	2007070632	A2	6/2007
WO	2008011194	A2	1/2008
WO	WO 2008025791	A1 *	3/2008

OTHER PUBLICATIONS

Repka et al., "Bioadhesive Properties of hydroxypropylcellulose topical films produced by hot melt extrusion," Journal of Controlled Release, 70: 341-351 (2001).

Repka et al., "Influence of Vitamin E TPGS on the properties of hydrophilic films produced by hot melt extrusion", International Journal of Pharmaceutics 202: 63-70 (2000).

Peh and Wong, Polymeric Films as Vehicle for Buccal Delivery: Swelling, Mechanical, and Bioadhesive Properties, J Pharm Pharmaceut Sci (www.ualberta.ca/~csps) 2 (2):53-61, 1999.

Bodmeier. Pharmaceutical Research, vol. 6, No. 8, 1989.

"Suboxone Subligualtabletten" In: Verlag Rote Liste Service GmbH: "Rote Liste 2008" 2008, Verlag Rote Liste Service GmbH, Frankfurt/Main, XP002624986, p. 39018, the whole document.

Notification of Transmittal of the International Search Report and the Written Opinion of the International Searching Authority, or the Declaration for International Application No. PCT/US2010/044488 dated Apr. 11, 2011.

Written Opinion of the International Searing Authority for International Application No. PCT/US2010/044488 dated Apr. 11, 2011.

Abeer M. Al-Ghananeem et al., "Effect of pH on Sublingual Absorption of Oxycodone Hydrochloride." AAPS PharmSciTech 2006; 7(1) Article 23 (http://www.aapspharmscitech.org).

Mahmood et al., "A limited sampling method for the estimation of AUC and Cmax of carbamazepine and carbamazepine epoxide following a single and multiple dose of a sustained-release product." BrJ Clin Pharmacol 1998; 45: pp. 241-246.

* cited by examiner

SUBLINGUAL AND BUCCAL FILM COMPOSITIONS

FIELD OF THE INVENTION

The present invention relates to compositions, methods of manufacture, products and methods of use relating to films containing therapeutic actives. The invention more particularly relates to self-supporting film dosage forms which provide a therapeutically effective dosage, essentially matching that of currently-marketed tablets containing the same active. Such compositions are particularly useful for treating narcotic dependence while providing sufficient buccal adhesion of the dosage form.

BACKGROUND OF THE RELATED TECHNOLOGY

Oral administration of two therapeutic actives in a single dosage form can be complex if the intention is to have one active absorbed into the body and the other active remain substantially unabsorbed. For example, one active may be relatively soluble in the mouth at one pH, and the other active may be relatively insoluble at the same pH. Moreover, the absorption kinetics of each therapeutic agent may be substantially different due to differing absorption of the charged and uncharged species. These factors represent some of the challenges in appropriately co-administering therapeutic agents.

Co-administration of therapeutic agents has many applications. Among such areas of treatment include treating individuals who suffer from narcotic dependence. Such individuals have a tendency to suffer from serious physical dependence on the narcotic, resulting in potentially dangerous withdrawal effects when the narcotic is not administered to the individual. In order to help individuals addicted to narcotics, it is known to provide a reduced level of a drug, which provides an effect of satisfying the body's urge for the narcotic, but does not provide the "high" that is provided by the misuse of the narcotic. The drug provided may be an agonist or a partial agonist, which provides a reduced sensation and may help lower dependence on the drug. However, even though these drugs provide only a low level of euphoric effect, they are capable of being abused by the individuals parenterally. In such cases, it is desirable to provide a combination of the drug with a second drug, which may decrease the likelihood of diversion and abuse of the first drug. For example, it is known to provide a dosage of an antagonist in combination with the agonist or partial agonist. The narcotic antagonist binds to a receptor in the brain to block the receptor, thus reducing the effect of the agonist.

One such combination of drugs has been marketed under the trade name Suboxone® as an orally ingestible tablet. However, such combinations in tablet form have the potential for abuse. In some instances, the patient who has been provided the drug may store the tablet in his mouth without swallowing the tablet, then later extract the agonist from the tablet and inject the drug into an individual's body. Although certain antagonists (such as highly water-soluble antagonists) may be used to help reduce the ability to separate the agonist, the potential for abuse still exists. It is desired to provide a dosage that cannot be easily removed from the mouth once it has been administered.

There is currently a need for an orally dissolvable film dosage form that provides the desired absorption levels of the agonist and antagonist, while providing an adhesive effect in the mouth, rendering it difficult to remove once placed in the mouth, thereby making abuse of the agonist difficult.

SUMMARY OF THE INVENTION

In one embodiment of the present invention, there is provided a film dosage composition including: a polymeric carrier matrix; a therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof, a therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof; and a buffer in an amount to provide a pH of the composition of a value sufficient to optimize absorption of the buprenorphine.

In another embodiment of the present invention, there is provided a film dosage composition including: a polymeric carrier matrix; a therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof, a therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof; and a buffer in an amount sufficient to inhibit the absorption of the naloxone when administered orally.

In still other embodiments, there may be provided a film dosage composition including: a polymeric carrier matrix; a therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof, a therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof; and a buffering system; where the buffering system includes a buffer capacity sufficient to maintain the ionization of naloxone during the time which the composition is in the oral cavity of a user.

In another embodiment of the invention, there is provided a method of treating narcotic dependence of a user, including the steps of: providing a composition including: a polymeric carrier matrix; a therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof, a therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof, and a buffer in an amount to provide a pH of the composition of a value sufficient to optimize absorption of the buprenorphine; and administering the composition to the oral cavity of a user.

In still another embodiment of the invention, there is provided a process of forming a film dosage composition including the steps of: casting a film-forming composition, the film-forming composition including: a polymeric carrier matrix; a therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof, a therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof, and a buffer in an amount to provide a pH of the composition of a value sufficient to optimize absorption of the buprenorphine and drying the film-forming composition to form a self-supporting film dosage composition.

In another embodiment, there is provided a film dosage composition including a therapeutically sufficient amount of buprenorphine or a pharmaceutically acceptable salt thereof and a therapeutically sufficient amount of naloxone or a pharmaceutically acceptable salt thereof, the film dosage composition having a bioequivalent release profile as compared to a Suboxone® tablet containing about 2 times the amount of buprenorphine or a pharmaceutically acceptable salt thereof.

Still other embodiments of the present invention provide an orally dissolving film formulation including buprenorphine and naloxone, where the formulation provides an in-vivo plasma profile having a Cmax of between about 0.624 ng/ml and about 5.638 ng/ml for buprenorphine and an in-vivo

plasma profile having a Cmax of between about 41.04 pg/ml to about 323.75 pg/ml for naloxone.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Definitions

As used herein, the term Cmax refers to the mean maximum plasma concentration after administration of the composition to a human subject. As also used herein, the term AUC refers to the mean area under the plasma concentration-time curve value after administration of the compositions formed herein. As will be set forth in more detail below, the term "optimizing the absorption" does not refer to reaching the maximum absorption of the composition, and rather refers to reaching the optimum level of absorption at a pH of about 2 to about 4. The "optimum" absorption may be, for example, a level that provides a bioequivalent absorption as administration of the currently available Suboxone® tablet. An "optimum" Cmax of buprenorphine is about 0.67 to about 5.36 mg/ml at dosages of from 2-16 mg buprenorphine at a given pH. Similarly, an "optimum" AUC of buprenorphine may be about 7.43 to about 59.46 hr*ng/ml at dosages of from 2-16 mg buprenorphine at a given pH. As will be described in more detail below, it has been surprisingly discovered that the absorption of one particular agonist, buprenorphine, can provide an optimum absorption at a pH of about 2-4 as well as about 5.5-6.5. Thus, one may "optimize" the absorption of buprenorphine by providing a pH of about 2-4 or about 5.5-6.5.

"Maximizing the absorption" refers to the maximum in vivo absorption values achieved at a pH of about 4 to about 9.

The term "local pH" refers to the pH of the region of the carrier matrix immediately surrounding the active agent as the matrix hydrates and/or dissolves, for example, in the mouth of the user.

By "inhibiting" the absorption of an active, it is meant achieving as complete an ionization state of the active as possible, such that little to none of the active is measurably absorbable. For example, at a pH of 3-3.5, the Cmax of an active such as naloxone for dosage of 0.5 mg to 4.0 mg ranges from 32.5 to 260 pg/ml, and an AUC of naloxone for dosage of 0.5 mg to 4.0 mg ranges from 90.55 to 724.4 hr*pg/ml. It is understood that at a pH lower than 3.0, further ionization would be expected and thus result in lower absorption.

The term "bioequivalent" means obtaining 80% to 125% of the Cmax and AUC values for a given active in a different product. For example, assuming Cmax and AUC values of buprenorphine for a commercially-available Suboxone® tablet (containing 2 mg buprenorphine and 0.5 mg naloxone) are 0.780 ng/ml and 6.789 hr*ng/ml, respectively, a bioequivalent product would have a Cmax of buprenorphine in the range of 0.624-0.975 ng/ml, and an AUC value of buprenorphine of 5.431-8.486 hr*ng/ml.

It will be understood that the term "film" includes thin films and sheets, in any shape, including rectangular, square, or other desired shape. The films described herein may be any desired thickness and size such that it may be placed into the oral cavity of the user. For example, the films may have a relatively thin thickness of from about 0.1 to about 10 mils, or they may have a somewhat thicker thickness of from about 10 to about 30 mils. For some films, the thickness may be even larger, i.e., greater than about 30 mils. Films may be in a single layer or they may be multi-layered, including laminated films.

Oral dissolving films generally fall into three main classes: fast dissolving, moderate dissolving and slow dissolving. Fast dissolving films generally dissolve in about 1 second to about 30 seconds in the mouth. Moderate dissolving films generally dissolve in about 1 to about 30 minutes in the mouth, and slow dissolving films generally dissolve in more than 30 minutes in the mouth. Fast dissolving films may consist of low molecular weight hydrophilic polymers (i.e., polymers having a molecular weight between about 1,000 to 9,000, or polymers having a molecular weight up to 200,000). In contrast, slow dissolving films generally have high molecular weight polymers (i.e., having a molecular weight in the millions).

Moderate dissolving films tend to fall in between the fast and slow dissolving films. Moderate dissolving films dissolve rather quickly, but also have a good level of mucoadhesion. Moderate dissolving films are also flexible, quickly wettable, and are typically non-irritating to the user. For the instant invention, it is preferable to use films that fall between the categories of fast dissolving and moderate dissolving. Such moderate dissolving films provide a quick enough dissolution rate, most desirably between about 1 minute and about 20 minutes, while providing an acceptable mucoadhesion level such that the film is not easily removable once it is placed in the oral cavity of the user.

Inventive films described herein may include one or more agonists or partial agonists used for the treatment of drug addiction. As used herein, the term "agonist" refers to a chemical substance that is capable of providing a physiological response or activity in the body of the user. The films described herein may further include one or more antagonists. As used herein, the term "antagonist" refers to any chemical substance that acts within the body of the user to reduce the physiological activity of another chemical substance. In some embodiments, an antagonist used herein may act to reduce and/or block the physiological activity of the agonist. The actives may be water-soluble, or they may be water-insoluble. As used herein, the term "water-soluble" refers to substances that are at least partially dissolvable in a solvent, including but not limited to water. The term "water-soluble" does not necessarily mean that the substance is 100% dissolvable in the solvent. The term "water-insoluble" refers to substances that are not dissolvable in a solvent, including but not limited to water. Solvents may include water, or alternatively may include other polar solvents by themselves or in combination with water.

Inventive Films

The present invention relates to methods of treating narcotic dependence in an individual. More desirably, the invention relates to the treatment of opioid dependence in an individual, while using a formulation and delivery that hinders misuse of the narcotic. Currently, treatment of opioid dependence is aided by administration of Suboxone®, which is an orally dissolvable tablet. This tablet which provides a combination of buprenorphine (an opioid agonist) and naloxone (an opioid antagonist). Therefore, the present invention provides a method of treating narcotic dependence by providing an orally dissolvable film dosage, which provides a bioequivalent effect to Suboxone®. The film dosage preferably provides buccal adhesion while it is in the user's mouth, rendering it difficult to remove after placement.

The film dosage composition preferably includes a polymeric carrier matrix. Any desired polymeric carrier matrix may be used, provided that it is orally dissolvable. Desirably, the dosage should have enough bioadhesion to not be easily removed and it should form a gel like structure when administered. The orally consumable films are preferably moderate-dissolving in the oral cavity and particularly suitable for

delivery of actives, although both fast and sustained release compositions are also among the various embodiments contemplated.

The films used in the pharmaceutical products may be produced by a combination of at least one polymer and a solvent, optionally including other fillers known in the art. The solvent may be water, a polar organic solvent including, but not limited to, ethanol, isopropanol, acetone, or any combination thereof. In some embodiments, the solvent may be a non-polar organic solvent, such as methylene chloride. The film may be prepared by utilizing a selected casting or deposition method and a controlled drying process. For example, the film may be prepared through controlled drying processes, which include application of heat and/or radiation energy to the wet film matrix to form a visco-elastic structure, thereby controlling the uniformity of content of the film. Such processes are described in more detail in commonly assigned U.S. application Ser. No. 10/074,272, filed on Feb. 14, 2002, and published as U.S. Patent Publication No. 2003/0107149 A1, the contents of which are incorporated herein by reference in their entirety. Alternatively, the films may be extruded as described in commonly assigned U.S. application Ser. No. 10/856,176, filed on May 28, 2004, and published as U.S. Patent Publication No. 2005/0037055 A1, the contents of which are incorporated herein by reference in their entirety.

The polymer included in the films may be water-soluble, water-swellable, water-insoluble, or a combination of one or more either water-soluble, water-swellable or water-insoluble polymers. The polymer may include cellulose or a cellulose derivative. Specific examples of useful water-soluble polymers include, but are not limited to, polyethylene oxide, pullulan, hydroxypropylmethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, polyvinyl pyrrolidone, carboxymethyl cellulose, polyvinyl alcohol, sodium alginate, polyethylene glycol, xanthan gum, tragancanth gum, guar gum, acacia gum, arabic gum, polyacrylic acid, methylmethacrylate copolymer, carboxyvinyl copolymers, starch, gelatin, and combinations thereof. Specific examples of useful water-insoluble polymers include, but are not limited to, ethyl cellulose, hydroxypropyl ethyl cellulose, cellulose acetate phthalate, hydroxypropyl methyl cellulose phthalate and combinations thereof. For higher dosages, it may be desirable to incorporate a polymer that provides a high level of viscosity as compared to lower dosages.

As used herein the phrase "water-soluble polymer" and variants thereof refer to a polymer that is at least partially soluble in water, and desirably fully or predominantly soluble in water, or absorbs water. Polymers that absorb water are often referred to as being water-swellable polymers. The materials useful with the present invention may be water-soluble or water-swellable at room temperature and other temperatures, such as temperatures exceeding room temperature. Moreover, the materials may be water-soluble or water-swellable at pressures less than atmospheric pressure. Desirably, the water-soluble polymers are water-soluble or water-swellable having at least 20 percent by weight water uptake. Water-swellable polymers having a 25 or greater percent by weight water uptake are also useful. In some embodiments, films formed from such water-soluble polymers may be sufficiently water-soluble to be dissolvable upon contact with bodily fluids.

Other polymers useful for incorporation into the films include biodegradable polymers, copolymers, block polymers and combinations thereof. It is understood that the term "biodegradable" is intended to include materials that chemically degrade, as opposed to materials that physically break apart (i.e., bioerodable materials). Among the known useful polymers or polymer classes which meet the above criteria are: poly(glycolic acid) (PGA), poly(lactic acid) (PLA), polydioxanes, polyoxalates, poly(α-esters), polyanhydrides, polyacetates, polycaprolactones, poly(orthoesters), polyamino acids, polyaminocarbonates, polyurethanes, polycarbonates, polyamides, poly(alkyl cyanoacrylates), and mixtures and copolymers thereof. Additional useful polymers include, stereopolymers of L- and D-lactic acid, copolymers of bis(p-carboxyphenoxy)propane acid and sebacic acid, sebacic acid copolymers, copolymers of caprolactone, poly (lactic acid)/poly(glycolic acid)/polyethyleneglycol copolymers, copolymers of polyurethane and (poly(lactic acid), copolymers of polyurethane and poly(lactic acid), copolymers of α-amino acids, copolymers of α-amino acids and caproic acid, copolymers of α-benzyl glutamate and polyethylene glycol, copolymers of succinate and poly(glycols), polyphosphazene, polyhydroxy-alkanoates and mixtures thereof. Binary and ternary systems are contemplated.

Other specific polymers useful include those marketed under the Medisorb and Biodel trademarks. The Medisorb materials are marketed by the Dupont Company of Wilmington, Del. and are generically identified as a "lactide/glycolide co-polymer" containing "propanoic acid, 2-hydroxy-polymer with hydroxy-polymer with hydroxyacetic acid." Four such polymers include lactide/glycolide 100 L, believed to be 100% lactide having a melting point within the range of 338°-347° F. (170°-175° C.); lactide/glycolide 100 L, believed to be 100% glycolide having a melting point within the range of 437°-455° F. (225°-235° C.); lactide/glycolide 85/15, believed to be 85% lactide and 15% glycolide with a melting point within the range of 338°-347° F. (170°-175° C.); and lactide/glycolide 50/50, believed to be a copolymer of 50% lactide and 50% glycolide with a melting point within the range of 338°-347° F. (170°-175° C.).

The Biodel materials represent a family of various polyanhydrides which differ chemically.

Although a variety of different polymers may be used, it is desired to select polymers that provide mucoadhesive properties to the film, as well as a desired dissolution and/or disintegration rate. In particular, the time period for which it is desired to maintain the film in contact with the mucosal tissue depends on the type of active contained in the composition. Some actives may only require a few minutes for delivery through the mucosal tissue, whereas other actives may require up to several hours or even longer. Accordingly, in some embodiments, one or more water-soluble polymers, as described above, may be used to form the film. In other embodiments, however, it may be desirable to use combinations of water-soluble polymers and polymers that are water-swellable, water-insoluble and/or biodegradable, as provided above. The inclusion of one or more polymers that are water-swellable, water-insoluble and/or biodegradable may provide films with slower dissolution or disintegration rates than films formed from water-soluble polymers alone. As such, the film may adhere to the mucosal tissue for longer periods or time, such as up to several hours, which may be desirable for delivery of certain active components.

Desirably, the individual film dosage has a small size, which is between about 0.5-1 inch by about 0.5-1 inch. Most preferably, the film dosage is about 0.75 inches×0.5 inches. The film dosage should have good adhesion when placed in the buccal cavity or in the sublingual region of the user. Further, the film dosage should disperse and dissolve at a moderate rate, most desirably dispersing within about 1 minute and dissolving within about 3 minutes. In some

embodiments the film dosage may be capable of dispersing and dissolving at a rate of between about 1 to about 1.5 minutes.

For instance, in some embodiments, the films may include polyethylene oxide alone or in combination with a second polymer component. The second polymer may be another water-soluble polymer, a water-swellable polymer, a water-insoluble polymer, a biodegradable polymer or any combination thereof. Suitable water-soluble polymers include, without limitation, any of those provided above. In some embodiments, the water-soluble polymer may include hydrophilic cellulosic polymers, such as hydroxypropyl cellulose and/or hydroxypropylmethyl cellulose. In accordance with some embodiments, polyethylene oxide may range from about 20% to 100% by weight in the polymer component, more specifically about 30% to about 70% by weight, and even more specifically about 40% to about 60% by weight. In some embodiments, one or more water-swellable, water-insoluble and/or biodegradable polymers also may be included in the polyethylene oxide-based film. Any of the water-swellable, water-insoluble or biodegradable polymers provided above may be employed. The second polymer component may be employed in amounts of about 0% to about 80% by weight in the polymer component, more specifically about 30% to about 70% by weight, and even more specifically about 40% to about 60% by weight.

The molecular weight of the polyethylene oxide also may be varied. In some embodiments, high molecular weight polyethylene oxide, such as about 4 million, may be desired to increase mucoadhesivity of the film. In some other embodiments, the molecular weight may range from about 100,000 to 900,000, more specifically from about 100,000 to 600,000, and even more specifically from about 100,000 to 300,000. In some embodiments, it may be desirable to combine high molecular weight (600,000 to 900,000) with low molecular weight (100,000 to 300,000) polyethylene oxide in the polymer component.

A variety of optional components and fillers also may be added to the films. These may include, without limitation: surfactants; plasticizers; polyalcohols; anti-foaming agents, such as silicone-containing compounds, which promote a smoother film surface by releasing oxygen from the film; thermo-setting gels such as pectin, carageenan, and gelatin, which help in maintaining the dispersion of components; inclusion compounds, such as cyclodextrins and caged molecules; coloring agents; and flavors. In some embodiments, more than one active components may be included in the film.

Additives may be included in the films. Examples of classes of additives include excipients, lubricants, buffering agents, stabilizers, blowing agents, pigments, coloring agents, fillers, bulking agents, sweetening agents, flavoring agents, fragrances, release modifiers, adjuvants, plasticizers, flow accelerators, mold release agents, polyols, granulating agents, diluents, binders, buffers, absorbents, glidants, adhesives, anti-adherents, acidulants, softeners, resins, demulcents, solvents, surfactants, emulsifiers, elastomers and mixtures thereof. These additives may be added with the active ingredient(s).

Useful additives include, for example, gelatin, vegetable proteins such as sunflower protein, soybean proteins, cotton seed proteins, peanut proteins, grape seed proteins, whey proteins, whey protein isolates, blood proteins, egg proteins, acrylated proteins, water-soluble polysaccharides such as alginates, carrageenans, guar gum, agar-agar, xanthan gum, gellan gum, gum arabic and related gums (gum ghatti, gum karaya, gum tragancanth), pectin, water-soluble derivatives of cellulose: alkylcelluloses hydroxyalkylcelluloses and hydroxyalkylalkylcelluloses, such as methylcelulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxyethylmethylcellulose, hydroxypropylmethylcellulose, hydroxybutylmethylcellulose, cellulose esters and hydroxyalkylcellulose esters such as cellulose acetate phthalate (CAP), hydroxypropylmethylcellulose (HPMC); carboxyalkylcelluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters such as carboxymethylcellulose and their alkali metal salts; water-soluble synthetic polymers such as polyacrylic acids and polyacrylic acid esters, polymethacrylic acids and polymethacrylic acid esters, polyvinylacetates, polyvinylalcohols, polyvinylacetatephthalates (PVAP), polyvinylpyrrolidone (PVP), PVY/vinyl acetate copolymer, and polycrotonic acids; also suitable are phthalated gelatin, gelatin succinate, crosslinked gelatin, shellac, water-soluble chemical derivatives of starch, cationically modified acrylates and methacrylates possessing, for example, a tertiary or quaternary amino group, such as the diethylaminoethyl group, which may be quaternized if desired; and other similar polymers.

Such extenders may optionally be added in any desired amount desirably within the range of up to about 80%, desirably about 3% to 50% and more desirably within the range of 3% to 20% based on the weight of all film components.

Further additives may flow agents and opacifiers, such as the oxides of magnesium aluminum, silicon, titanium, etc. desirably in a concentration range of about 0.02% to about 3% by weight and desirably about 0.02% to about 1% based on the weight of all film components.

Further examples of additives are plasticizers which include polyalkylene oxides, such as polyethylene glycols, polypropylene glycols, polyethylene-propylene glycols, organic plasticizers with low molecular weights, such as glycerol, glycerol monoacetate, diacetate or triacetate, triacetin, polysorbate, cetyl alcohol, propylene glycol, sorbitol, sodium diethylsulfosuccinate, triethyl citrate, tributyl citrate, and the like, added in concentrations ranging from about 0.5% to about 30%, and desirably ranging from about 0.5% to about 20% based on the weight of the polymer.

There may further be added compounds to improve the texture properties of the starch material such as animal or vegetable fats, desirably in their hydrogenated form, especially those which are solid at room temperature. These fats desirably have a melting point of 50° C. or higher. Preferred are tri-glycerides with C_{12}-, C_{14}-, C_{16}-, C_{18}-, C_{20}- and C_{22}-fatty acids. These fats can be added alone without adding extenders or plasticizers and can be advantageously added alone or together with mono- and/or di-glycerides or phosphatides, especially lecithin. The mono- and di-glycerides are desirably derived from the types of fats described above, i.e. with C_{12}-, C_{14}-, C_{16}-, C_{18}-, C_{20}- and C_{22}-fatty acids.

The total amounts used of the fats, mono-, di-glycerides and/or lecithins are up to about 5% and preferably within the range of about 0.5% to about 2% by weight of the total film composition.

It further may be useful to add silicon dioxide, calcium silicate, or titanium dioxide in a concentration of about 0.02% to about 1% by weight of the total composition. These compounds act as flow agents and opacifiers.

Lecithin is one surface active agent for use in the films described herein. Lecithin may be included in the feedstock in an amount of from about 0.25% to about 2.00% by weight. Other surface active agents, i.e. surfactants, include, but are not limited to, cetyl alcohol, sodium lauryl sulfate, the Spans™ and Tweens™ which are commercially available from ICI Americas, Inc. Ethoxylated oils, including ethoxylated castor oils, such as Cremophor EL which is commer-

cially available from BASF, are also useful. Carbowax™ is yet another modifier which is very useful in the present invention. Tweens™ or combinations of surface active agents may be used to achieve the desired hydrophilic-lipophilic balance ("HLB"). The present invention, however, does not require the use of a surfactant and films or film-forming compositions of the present invention may be essentially free of a surfactant while still providing the desirable uniformity features of the present invention.

Other ingredients include binders which contribute to the ease of formation and general quality of the films. Non-limiting examples of binders include starches, pregelatinize starches, gelatin, polyvinylpyrrolidone, methylcellulose, sodium carboxymethylcellulose, ethylcellulose, polyacrylamides, polyvinyloxoazolidone, and polyvinylalcohols.

Further potential additives include solubility enhancing agents, such as substances that form inclusion compounds with active components. Such agents may be useful in improving the properties of very insoluble and/or unstable actives. In general, these substances are doughnut-shaped molecules with hydrophobic internal cavities and hydrophilic exteriors. Insoluble and/or instable actives may fit within the hydrophobic cavity, thereby producing an inclusion complex, which is soluble in water. Accordingly, the formation of the inclusion complex permits very insoluble and/or instable actives to be dissolved in water. A particularly desirable example of such agents are cyclodextrins, which are cyclic carbohydrates derived from starch. Other similar substances, however, are considered well within the scope of the present invention.

Suitable coloring agents include food, drug and cosmetic colors (FD&C), drug and cosmetic colors (D&C), or external drug and cosmetic colors (Ext. D&C). These colors are dyes, their corresponding lakes, and certain natural and derived colorants. Lakes are dyes absorbed on aluminum hydroxide.

Other examples of coloring agents include known azo dyes, organic or inorganic pigments, or coloring agents of natural origin. Inorganic pigments are preferred, such as the oxides or iron or titanium, these oxides, being added in concentrations ranging from about 0.001 to about 10%, and preferably about 0.5 to about 3%, based on the weight of all the components.

Flavors may be chosen from natural and synthetic flavoring liquids. An illustrative list of such agents includes volatile oils, synthetic flavor oils, flavoring aromatics, oils, liquids, oleoresins or extracts derived from plants, leaves, flowers, fruits, stems and combinations thereof. A non-limiting representative list of examples includes mint oils, cocoa, and citrus oils such as lemon, orange, grape, lime and grapefruit and fruit essences including apple, pear, peach, grape, strawberry, raspberry, cherry, plum, pineapple, apricot or other fruit flavors.

Other useful flavorings include aldehydes and esters such as benzaldehyde (cherry, almond), citral i.e., alphacitral (lemon, lime), neral, i.e., beta-citral (lemon, lime), decanal (orange, lemon), aldehyde C-8 (citrus fruits), aldehyde C-9 (citrus fruits), aldehyde C-12 (citrus fruits), tolyl aldehyde (cherry, almond), 2,6-dimethyloctanol (green fruit), and 2-dodecenal (citrus, mandarin), combinations thereof and the like.

The sweeteners may be chosen from the following non-limiting list: glucose (corn syrup), dextrose, invert sugar, fructose, and combinations thereof, saccharin and its various salts such as the sodium salt; dipeptide sweeteners such as aspartame; dihydrochalcone compounds, glycyrrhizin; Stevia Rebaudiana (Stevioside); chloro derivatives of sucrose such as sucralose; sugar alcohols such as sorbitol, mannitol, xylitol, and the like. Also contemplated are hydrogenated starch hydrolysates and the synthetic sweetener 3,6-dihydro-6-methyl-1-1-1,2,3-oxathiazin-4-one-2,2-dioxide, particularly the potassium salt (acesulfame-K), and sodium and calcium salts thereof, and natural intensive sweeteners, such as Lo Han Kuo. Other sweeteners may also be used.

Anti-foaming and/or de-foaming components may also be used with the films. These components aid in the removal of air, such as entrapped air, from the film-forming compositions. Such entrapped air may lead to non-uniform films. Simethicone is one particularly useful anti-foaming and/or de-foaming agent. The present invention, however, is not so limited and other anti-foam and/or de-foaming agents may suitable be used.

As a related matter, simethicone and related agents may be employed for densification purposes. More specifically, such agents may facilitate the removal of voids, air, moisture, and similar undesired components, thereby providing denser, and thus more uniform films. Agents or components which perform this function can be referred to as densification or densifying agents. As described above, entrapped air or undesired components may lead to non-uniform films.

Simethicone is generally used in the medical field as a treatment for gas or colic in babies. Simethicone is a mixture of fully methylated linear siloxane polymers containing repeating units of polydimethylsiloxane which is stabilized with trimethylsiloxy end-blocking unites, and silicon dioxide. It usually contains 90.5-99% polymethylsiloxane and 4-7% silicon dioxide. The mixture is a gray, translucent, viscous fluid which is insoluble in water.

When dispersed in water, simethicone will spread across the surface, forming a thin film of low surface tension. In this way, simethicone reduces the surface tension of bubbles air located in the solution, such as foam bubbles, causing their collapse. The function of simethicone mimics the dual action of oil and alcohol in water. For example, in an oily solution any trapped air bubbles will ascend to the surface and dissipate more quickly and easily, because an oily liquid has a lighter density compared to a water solution. On the other hand, an alcohol/water mixture is known to lower water density as well as lower the water's surface tension. So, any air bubbles trapped inside this mixture solution will also be easily dissipated. Simethicone solution provides both of these advantages. It lowers the surface energy of any air bubbles that trapped inside the aqueous solution, as well as lowering the surface tension of the aqueous solution. As the result of this unique functionality, simethicone has an excellent anti-foaming property that can be used for physiological processes (anti-gas in stomach) as well as any for external processes that require the removal of air bubbles from a product.

In order to prevent the formation of air bubbles in the films, the mixing step can be performed under vacuum. However, as soon as the mixing step is completed, and the film solution is returned to the normal atmosphere condition, air will be re-introduced into or contacted with the mixture. In many cases, tiny air bubbles will be again trapped inside this polymeric viscous solution. The incorporation of simethicone into the film-forming composition either substantially reduces or eliminates the formation of air bubbles.

Simethicone may be added to the film-forming mixture as an anti-foaming agent in an amount from about 0.01 weight percent to about 5.0 weight percent, more desirably from about 0.05 weight percent to about 2.5 weight percent, and most desirably from about 0.1 weight percent to about 1.0 weight percent.

Any other optional components described in commonly assigned U.S. Pat. No. 7,425,292 and U.S. application Ser. No. 10/856,176, referred to above, also may be included in the films described herein.

When the dosage form includes at least one antagonist, it may be desired to control the release of the antagonist, so as to delay or wholly prevent the release of the antagonist from the dosage when taken orally. Desirably, the dosage form is a self-supporting film composition, which is placed into the oral cavity of the user. In a dosage form that is to be placed in the oral cavity, it is desired to absorb the agonist buccally, so as to provide rapid integration of the agonist into the body of the user. At the same time, it may be desired to prevent or reduce absorption of any antagonist buccally, thereby allowing the antagonist to be swallowed and destroyed in the stomach. Reducing the absorption of an antagonist may be achieved via physical means, such as by encapsulating the antagonist in a material that blocks absorption. It is desired, however, to reduce the absorption of the antagonist by chemical means, such as by controlling the local pH of the dosage.

It has been found that by controlling the local pH of the dosage form, the release and/or absorption of the actives therein may be controlled. For example, in a dosage that includes an amount of an agonist, the local pH may be controlled to a level that maximizes its release and/or absorption into the oral cavity of the user. In dosages incorporating an amount of an agonist and an amount of an antagonist, the local pH may be controlled to a level that maximizes the release and/or absorption of the agonist while simultaneously minimizing the release and/or absorption of the antagonist.

The dosage form preferably includes a combination of a partial agonist and an antagonist, while the dosage has a controlled pH. In one embodiment, the partial agonist may include buprenorphine or a pharmaceutically acceptable salt thereof, while the antagonist includes naloxone or a therapeutically acceptable salt thereof. It should be understood that the present invention is not limited to the use of buprenorphine and naloxone, and any agonist (or partial agonist) and any antagonist may be incorporated into the present invention for use in treatment of drug addiction. The agonist and optional antagonist should be selected from those agonists and antagonists that are useful in treating the particular narcotic dependence being treated.

As discussed above, the local pH of the dosage is preferably controlled to provide the desired release and/or absorption of the agonist and antagonist. Buprenorphine is known to have a pKa of about 8.42, while naloxone has a pKa of about 7.94. According to pH partition theory, one would expect that saliva (which has a pH of about 6.5) would maximize the absorption of both actives. However, it has been surprisingly discovered by the Applicants that by buffering the dosage to a particular pH level, the optimum levels of absorption of the agonist and antagonist may be achieved. Desirably, the local pH of a composition including an agonist and an antagonist is between about 2 to about 4, and most desirably is from 3 to 4. At this local pH level, the optimum absorption of the agonist and the antagonist is achieved. As will be described in more detail in the Examples below, controlling the local pH of the film compositions of the present invention provides a system in which the desired release and/or absorption of the components is bioequivalent to that of a similar Suboxone® tablet.

In one embodiment, the dosage form is a self-supporting film. In this embodiment, the film dosage includes a polymer carrier matrix, a therapeutically effective amount of buprenorphine, an agonist. The buffer is preferably capable of providing a local pH of the composition within a range that provides the desired level of absorption of the buprenorphine. The resulting dosage is a film composition that allows for a rapid and effective release of buprenorphine into the oral cavity of the user. At the same time, the film composition preferably has a sufficient adhesion profile, such that the film cannot easily be removed from the oral cavity of the user once it has been placed into the cavity. Full release of the buprenorphine preferably takes place within less than about thirty minutes, and preferably remains in the oral cavity for at least 1 minute.

As explained above, while providing a pharmaceutically acceptable level of an agonist is helpful in treating those with narcotic addiction, it may be desirable to provide the buprenorphine in combination with naloxone (an antagonist) so as to reduce the effect of the agonist and therefore aid in reducing dependency of the narcotic. Therefore, it may be desirable to combine the opioid agonist (or partial agonist) in the film composition with an opioid antagonist or a pharmaceutically acceptable salt thereof. The actives may be dispersed throughout the dosage separately or they may be combined together and dispersed into the dosage. Most desirably the antagonist includes naloxone, but any suitable basic antagonist may be selected as desired. The antagonist may optionally be water-soluble, so as to render separation of the antagonist and agonist difficult, thereby lessening the potential for abuse of the agonist.

As with a film including an agonist, the film including an agonist and an antagonist is desirably pH-controlled through the inclusion of a buffer. In such combination films, it has been discovered that the local pH of the film composition should preferably be in the range of about 2 to about 4, and more preferably about 3 to about 4 so as to provide a bioequivalent product as the commercially-available Suboxone® tablet. Most preferably the local pH of the film composition is about 3.5. At this local pH level, absorption of the buprenorphine is optimized while the absorption of the naloxone is inhibited.

The film may contain any desired level of self-supporting film forming polymer, such that a self-supporting film composition is provided. In one embodiment, the film composition contains a film forming polymer in an amount of at least 25% by weight of the composition. The film forming polymer may alternatively be present in an amount of at least 50% by weight of the composition. As explained above, any film forming polymers that impart the desired mucoadhesion and rate of film dissolution may be used as desired.

Any desired level of agonist and optional antagonist may be included in the dosage, so as to provide the desired effect. In one particular embodiment, the film composition includes about 2 mg to about 16 mg of agonist per dosage. More desirably, the film composition includes about 4 mg to about 12 mg of agonist per dosage. If desired, the film composition may include about 0.5 mg to about 5 mg of antagonist per dosage. More desirably, the film composition includes about 1 mg to about 3 mg of antagonist per dosage. If an antagonist is incorporated into the film, the film composition may include the antagonist in a ratio of about 6:1-2:1 agonist to antagonist. Most desirably, the film composition contains about 4:1 agonist to antagonist per dosage. For example, in one embodiment, the dosage includes an agonist in an amount of about 12 mg, and includes an antagonist in an amount of about 3 mg.

The film compositions further desirably contains a buffer so as to control the local pH of the film composition. Any desired level of buffer may be incorporated into the film composition so as to provide the desired local pH level. The buffer is preferably provided in an amount sufficient to control the release from the film and/or the absorption into the

body of the agonist and the optional antagonist. In a desired embodiment, the film composition includes buffer in a ratio of buffer to agonist in an amount of from about 2:1 to about 1:5 (buffer:agonist). The buffer may alternatively be provided in a 1:1 ratio of buffer to agonist. As stated above, the film composition preferably has a local pH of about 2 to about 4, and most preferably has a local pH of about 3.5. Any buffer system may be used as desired. In some embodiments, the buffer may include sodium citrate, citric acid, and combinations thereof.

In this embodiment, the resulting film composition includes a polymer matrix, an agonist, and an optional antagonist, while the film composition has a controlled local pH to the level desired. The buffer is preferably present in an amount to provide a therapeutically adequate absorption of the agonist, while simultaneously limiting the absorption of the antagonist. Controlling of the local pH allows for the desired release and/or absorption of the components, and thus provides a more useful and effective dosage.

The film dosage composition may include a polymer carrier matrix, a therapeutically effective amount of agonist, a therapeutically effective amount of antagonist, and a buffering system. The buffering system may include a buffer in addition to a solvent. The buffering system desirably includes a sufficient level of buffer so as to provide a desired local pH level of the film dosage composition.

In addition to a desired local pH level, the buffer preferably has a buffer capacity sufficient to maintain the ionization of the optional antagonist during the time that the composition is in the oral cavity of a user. Maintaining the ionization of the antagonist serves to limit the absorption of the antagonist, and thus provide the desired control of the antagonist. While the ionization of the antagonist is limited, the ionization of the agonist may not be so limited. As such, the resulting dosage form provides absorption of the agonist to the user, while sufficiently reducing and/or preventing absorption of the antagonist. By keeping the antagonist ionized and the local pH at the optimum pH, the antagonist has limited if any absorption, but is still present should the product be abused or taken via a different route of administration. However, when taken as administered, the antagonist has little to no effect in blocking the agonist.

The film dosage composition including an agonist may be configured to provide an in vivo plasma profile having a mean maximum plasma concentration (Cmax) in a desired range. It has been discovered by the Applicants that controlling the Cmax of the film composition allows one to control the absorption of the active (such as an agonist) into the user. The resulting film composition is more effective and suitable for delivery to a user.

As explained, the film dosage composition provides a bioequivalent result to a commercially available Suboxone® product. As will be explained more in the Examples below, commercially available Suboxone® provides different absorption levels depending on the amount of buprenorphine and naloxone administered. The present invention desirably provides a film product providing bioequivalent release as that of the Suboxone® product. As with the Suboxone® product, the buprenorphine may be present in an amount of from about 2 mg to about 16 mg per dosage, or, if desired about 4 mg to about 12 mg per dosage. Additionally, the naloxone may be present in any desired amount, preferably at about 25% the level of buprenorphine. For example, an inventive film product may have 2 mg buprenorphine and 0.5 mg naloxone, 4 mg buprenorphine and 1 mg naloxone, 8 mg buprenorphine and 2 mg naloxone, 12 mg buprenorphine and 3 mg naloxone, 16 mg buprenorphine and 4 mg naloxone, or any similar amounts.

It has further been discovered that, by controlling the mean area under the curve (AUC) value of the film composition, a more effective dosage form may be provided. As is described in more detail in the Examples below, the inventive film composition preferably provides an AUC value so as to provide a bioequivalent result as that provided by the commercially available Suboxone® tablet. In one embodiment, the film composition may include a mean AUCinf value of about 6.8 hr·ng/ml or greater. Alternatively, the film composition may include a mean AUCinf value of from about 6.8 hr·ng/ml to about 66 hr·ng/ml.

As explained above, the film compositions may include naloxone, an antagonist. When the film composition includes a combination of agonist and antagonist, the film composition may be configured to provide a particular Cmax and/or AUCinf for the antagonist. For example, when a buprenorphine agonist and a naloxone antagonist are incorporated into the film composition, the naloxone may be configured to provide a Cmax of less than about 400 pg/ml, less than about 318 pg/ml, less than about 235 pg/ml, less than about 92 pg/ml or less than about 64 pg/ml. In such films, the naloxone may provide a mean AUCinf value of less than about 1030 hr·ng/ml.

In formulations which include an agonist in combination with an antagonist, the film composition may be prepared to provide a desired Cmax and/or AUCinf value for each of the agonist and antagonist. In one embodiment, the film composition provides an in vivo plasma profile having a Cmax of less than about 6.4 ng/ml for the agonist and an in vivo plasma profile having a Cmax of less than about 400 pg/ml for the antagonist. In such embodiments, the formulation may provide an AUCinf value of more than about 6.8 hr·ng/ml for the agonist. If desired, the formulation may provide an AUCinf value of less than about 1030 hr·pg/ml for the antagonist. Such compositions may include the agonist and the antagonist in any desired amount, and in a preferred embodiment, the composition includes about 2 mg to about 16 mg of the agonist per dosage and about 0.5 mg to about 4 mg of the antagonist per dosage.

The present invention provides a method of treating narcotic dependence in a patient. In one embodiment, the patient is dependent on opioid narcotics, but the patient may have a dependence on non-opioid narcotics. Desirably, the patient is treated by providing a dosage to the patient, which provides an effective release of actives but simultaneously provides a suitable adhesion so that the dosage cannot be easily removed. In one method of treatment, an orally dissolvable film composition is provided to a patient.

Depending on the particular narcotic that the patient experiences dependence upon, the film composition may include one or more particular active components. In one embodiment, the film composition includes a polymer carrier matrix and a therapeutically effective amount of an agonist. Desirably the agonist is a partial agonist. For opioid dependency, the agonist may be an opioid agonist, such as buprenorphine or a pharmaceutically acceptable salt thereof. The film composition preferably includes a buffer in an amount sufficient to control the local pH of the film composition. Any buffer system may be used, including sodium citrate, citric acid, and combinations thereof. In compositions solely including an agonist, the local pH of the film composition is desirably about 5 to about 6.5, and most desirably the local pH is about 5.5. At this level, the absorption of the agonist is most effec-

tive. To treat the dependency, the film composition is administered to the patient, most desirably into the oral cavity of the patient.

If desired, the composition may include a therapeutically effective amount of an antagonist, to prevent abuse of the agonist. A "therapeutically effective amount" of an antagonist is intended to refer to an amount of the antagonist that may be useful in diverting abuse of the agonist by a user. The antagonist may be any desired antagonist, and in one embodiment includes naloxone or a pharmaceutically acceptable salt thereof. The film composition is preferably administered to a patient through the oral cavity of the patient, but may be administered in any desired means. The orally dissolvable film composition is then allowed to dissolve in the oral cavity of the patient for a sufficient time so as to release the active(s) therein. In some embodiments, the film composition may remain in the oral cavity for at least 30 seconds, and in some embodiments may remain in the oral cavity for at least 1 minute. After the film composition is placed into the oral cavity of the patient, the film preferably becomes sufficiently adhered so as to render its removal difficult. After the film composition has been administered to the patient, the active(s) are sufficiently released from the composition and allowed to take effect on the patient.

The film compositions of the present invention may be formed via any desired process. Suitable processes are set forth in U.S. Pat. Nos. 7,425,292 and 7,357,891, the entire contents of which are incorporated by reference herein. In one embodiment, the film dosage composition is formed by first preparing a wet composition, the wet composition including a polymeric carrier matrix, a therapeutically effective amount of an agonist, and a buffer in an amount sufficient to control the local pH of the composition to a desired level. The wet composition is cast into a film and then sufficiently dried to form a self-supporting film composition. The wet composition may be cast into individual dosages, or it may be cast into a sheet, where the sheet is then cut into individual dosages. The agonist may be a partial agonist. If desired, the wet composition may include a therapeutically effective amount of an antagonist.

The agonist and the optional antagonist are preferably selected to treat a particular narcotic dependency. For opioid dependency, for example, the agonist may include buprenorphine or a pharmaceutically acceptable salt thereof, while the antagonist may include naloxone or a pharmaceutically acceptable salt thereof. The local pH of the film composition is desirably maintained at about 2 to about 4.

EXAMPLES

Example 1

Composition of Buprenorphine/Naloxone Films at Various Strengths

Film strips including a combination of buprenorphine and naloxone were prepared. Four different strength film compositions were prepared, which include a ratio of buprenorphine to naloxone of 16/4, 12/3, 8/2, and 2/0.5. The compositions are summarized in Table 1 below.

TABLE 1

Various Compositions of Film Dosages

Components	Buprenorphine/ Naloxone Films Unit Formula (mg per film strip) Buprenorphine/ Naloxone Ratios 16/4	12/3	8/2	2/0.5
Active Components				
Buprenorphine HCl	17.28	12.96	8.64	2.16
Naloxone HCl Dihydrate	4.88	3.66	2.44	0.61
Inactive Components				
Polyethylene Oxide, NF (MW 200,000)	27.09	20.32	13.55	—
Polyethylene Oxide, NF (MW 100,000)	12.04	9.03	6.02	19.06
Polyethylene Oxide, NF (MW 900,000)	4.82	3.62	2.41	2.05
Maltitol, NF	12.04	9.03	6.02	5.87
Flavor	6.0	4.5	3.0	2.4
Citric Acid, USP	5.92	4.44	2.96	2.96
HPMC	4.22	3.16	2.11	2.34
Ace-K	3.0	2.25	1.5	1.2
Sodium Citrate, anhydrous	2.68	2.01	1.34	1.34
Colorant	0.03	0.02	0.01	0.01
Total (mg)	100	75	50	40

Example 2

Absorption Studies for Suboxone® Products

Various film and tablet products were prepared and tested for absorption data, including Cmax and AUC absorption levels. The products tested included Suboxone® tablets made with either 2 mg or 16 mg buprenorphine as well as either 0.5 mg or 4.0 mg naloxone. For 16 mg buprenorphine tablets, two 8 mg buprenorphine tablets were combined together to provide the level of components of a 16 mg buprenorphine tablet. In instances where a 12 mg buprenorphine tablet was evaluated, this dosage was obtained by combining one 8 mg buprenorphine tablet and two 2 mg buprenorphine tablets. These products were tested for absorption levels, with the amounts listed in Table 2 below.

TABLE 2

Absorption Data for Suboxone ® products

Sample	C max	AUC
Buprenorphine (2 mg) Suboxone ® Tablet	0.780 ng/ml	6.789 hr * ng/ml
Naloxone (0.5 mg) Suboxone ® Tablet	51.30 pg/ml	128.60 hr * pg/ml
Buprenorphine (16 mg) Suboxone ® Tablet	4.51 ng/ml	44.99 hr * ng/ml
Naloxone (4 mg) Suboxone ® Tablet	259.00 pg/ml	649.60 hr * pg/ml

Using the data from Table 2, absorption data for the Suboxone® tablets for other levels of buprenorphine and naloxone are set forth in Table 2A below.

TABLE 2A

Extrapolated Absorption Data for Suboxone ® products

Sample	C max	AUC
Buprenorphine (4 mg) Suboxone ® Tablet	1.35 ng/ml	12.25 hr * ng/ml
Naloxone (1 mg) Suboxone ® Tablet	80.97 pg/ml	203 hr * pg/ml
Buprenorphine (8 mg) Suboxone ® Tablet	2.29 ng/ml	23.17 hr * ng/ml
Naloxone (2 mg) Suboxone ® Tablet	140.31 pg/ml	351.8 hr * pg/ml
Buprenorphine (12 mg) Suboxone ® Tablet	3.23 ng/ml	34.08 hr * ng/ml
Naloxone (3 mg) Suboxone ® Tablet	199.7 pg/ml	500.6 hr * pg/ml

Example 3

Evaluation of Bioequivalence of Suboxone® Tablets

Using the data generated for Suboxone® tablets in Table 2 above, acceptable bioequivalence ranges are generated to provide an equivalent treatment level as the Suboxone® tablet. As currently understood, a product provides a bioequivalent effect if it provides absorption levels between about 80% to about 125% of the Suboxone® tablet. Absorption in this range is considered to be bioequivalent.

TABLE 3

Acceptable Bioequivalence Ranges for Suboxone ® Tablets (80 to 125%)

Description of Sample	C max	AUC
Buprenorphine 2 mg	0.624 to 0.975 ng/ml	5.431 to 8.486 hr * ng/ml
Naloxone 0.5 mg	41.04 to 64.13 pg/ml	102.88 to 160.75 hr * pg/ml
Buprenorphine 16 mg	3.608 to 5.638 ng/ml	35.992 to 56.238 hr * ng/ml
Naloxone 4 mg	207.20 to 323.75 pg/ml	519.68 to 812.00 hr * pg/ml

Thus, to be considered bioequivalent to the Suboxone® tablet, the Cmax of buprenorphine is between about 0.624 and 5.638, and the AUC of buprenorphine is between about 5.431 to about 56.238. Similarly, to be considered bioequivalent to the Suboxone® tablet, the Cmax of naloxone is between about 41.04 to about 323.75, and the AUC of naloxone is between about 102.88 to about 812.00.

Example 4

Absorption Studies for Film Products at pH 3.5

Various film products were prepared and tested for absorption data, including Cmax and AUC absorption levels. The products tested included inventive film strips, the film strips having either 2 mg or 16 mg buprenorphine as well as either 0.5 mg or 4.0 mg naloxone. These products were tested for absorption levels, with the amounts listed in Table 4 below.

TABLE 4

Absorption Data for inventive film products at pH 3.5

Sample	C max	AUC
Buprenorphine (2 mg) Sublingual Film	0.947 ng/ml	7.82 hr * ng/ml
Naloxone (0.5 mg) Sublingual Film	51.10 pg/ml	128.60 hr * pg/ml

TABLE 4-continued

Absorption Data for inventive film products at pH 3.5

Sample	C max	AUC
Buprenorphine (16 mg) Sublingual Film	5.47 ng/ml	55.30 hr * ng/ml
Naloxone (4 mg) Sublingual Film	324.00 pg/ml	873.60 hr * pg/ml

As can be seen, in this experiment, the values for buprenorphine absorbance were squarely in the bioequivalence range evaluated above. The inventive films were therefore determined to have provided a bioequivalent absorption of buprenorphine at a local pH of 3.5 as the commercially available Suboxone® tablet. The values for absorption of naloxone were very close to the bioequivalent range of Suboxone®. The slightly higher absorption of Naloxone was not due to the local pH but rather to the amount of buffer (buffer capacity as discussed in the application). This is confirmed by the fact that the lower 2/0.5 mg dose is in range for the Naloxone and this is due to the higher buffer capacity for the 2/0.5 dose as pointed out in the buffer capacity chart.

Example 5

Preparation of Films for In Vivo Study

Film dosages were prepared for use in an in vivo study to determine the bioavailability of buprenorphine/naloxone tablets and film formulations. Specifically, the films were tested to determine whether the film provides a bioequivalent effect to that of a tablet formulation.

Three film formulations including 8 mg buprenorphine and 2 mg naloxone were prepared, each being buffered to a different pH. The first film did not include any buffer, providing a local pH of about 6.5. The second was buffered to a local pH level of about 3-3.5. The third was buffered to a local pH value of about 5-5.5. The formulations are set forth in Table 5 below.

TABLE 5

Formulations of Test Films at Various pH Levels

	Test formulation 1 8 mg/2 mg pH = 6.5		Test formulation 2 8 mg/2 mg pH = 3-3.5		Test formulation 3 8 mg/2 mg pH = 5-5.5	
Component	% w/w	Mg/film	% w/w	Mg/film	% w/w	Mg/film
Buprenorphine HCl	21.61	8.64	17.28	8.64	17.28	8.64
Naloxone HCl Dihydrate	6.10	2.44	4.88	2.44	4.88	2.44
Polymer	5.05	2.02	4.82	2.41	4.82	2.41
Polymer	28.48	11.39	27.09	13.55	27.09	13.55
Polymer	12.65	5.06	12.04	6.02	12.04	6.02
Polymer	4.43	1.77	4.22	2.11	4.22	2.11
Sweetener	12.65	5.06	12.04	6.02	12.04	6.02
Sweetener	3	1.2	3	1.5	3	1.5
Flavor	6	2.4	6	3	6	3
Citric acid	0	0	5.92	2.96	2.51	1.26
Sodium citrate	0	0	2.68	1.34	6.08	3.04
FD&C yellow #6	0.025	0.01	0.03	0.02	0.03	0.02
Total	100	40	100	50	100	50

Example 6

Analysis of In Vivo Absorption of Film Having a pH of 6.5

The film dosage composition of film having a local pH of 6.5 was analyzed. Specifically, Test Formulation 1, as prepared in Example 5 was analyzed in vivo to determine the absorption of buprenorphine and of naloxone. The comparative film was compared to the absorption of buprenorphine and of naloxone provided by a one dose tablet (Suboxone®). The test film was compared to determine whether it provided a bioequivalent effect as the tablet product.

The results for Test Formulation 1, which had a local pH of about 6.5, as compared to the one dose tablet, are set forth in Tables 6 and 7 below.

TABLE 6

Buprenorphine In Vivo Absorption Data for Test Formulation 1

	Suboxone ® sublingual				Test Formulation 1 (pH = 6.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
T_{max} (hr)	15	1.60	0.47	29.41	15	1.50	0.62	41.23
C_{max} (ng/mL)	15	2.27	0.562	24.77	15	2.60	0.872	33.53
AUC_{last} (hr * ng/mL)	15	27.08	10.40	38.41	15	31.00	12.93	41.72
AUC_{inf} (hr * ng/mL)	15	29.58	11.15	37.68	15	33.37	13.88	41.61
$T_{1/2}$ (hr)	15	44.76	20.86	46.60	15	40.73	14.93	36.66

TABLE 7

Naloxone In Vivo Absorption Data for Test Formulation 1

	Suboxone ® sublingual				Test Formulation 1 (pH = 6.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
T_{max} (hr)	15	0.90	0.23	25.32	15	0.68	0.18	25.75
C_{max} (pg/mL)	15	94.6	39.1	41.33	15	410	122	29.75
AUC_{last} (hr * pg/mL)	15	297.1	120.7	40.62	15	914.8	158.1	17.29
AUC_{inf} (hr * pg/mL)	15	306.1	122.6	40.06	15	924.2	158.8	17.18
$T_{1/2}$ (hr)	15	6.62	2.60	39.26	15	6.86	2.08	30.27

As can be seen, the in vivo data indicates that buprenorphine is absorbed very well from the film formulation at a local pH of 6.5, and matched closely the absorption seen in the Suboxone® one dose tablet. However, the absorption was also maximized for the naloxone, which was undesirable. It was determined that a film having a combination of buprenorphine and naloxone and a local pH of 6.5 did not provide a bioequivalent effect as the Suboxone® tablet for both buprenorphine and naloxone.

Example 7

Analysis of In Vivo Absorption of Film Having a pH of 5-5.5

Having determined the absorption of buprenorphine and naloxone in film having a local pH of 6.5, a film dosage composition of film having a local pH of 5-5.5 was analyzed. Specifically, Test Formulation 3, as prepared in Example 5 was analyzed in vivo to determine the absorption of buprenorphine and of naloxone. The comparative films were compared to the absorption of buprenorphine and of naloxone provided by the Suboxone® one dose tablet. The test film was compared to determine whether it provided a bioequivalent effect as the Suboxone® tablet.

The results for Test Formulation 3, which had a local pH of about 5-5.5, as compared to the Suboxone® tablet, are set forth in Tables 8 and 9 below.

TABLE 8

Buprenorphine In Vivo Absorption Data for Test Formulation 3

	Suboxone ® sublingual				Test Formulation 3 (pH = 5-5.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
T_{max} (hr)	15	1.60	0.47	29.41	14	1.50	0.43	28.50
C_{max} (ng/mL)	15	2.27	0.562	24.77	14	3.47	1.57	45.40
AUC_{last} (hr * ng/mL)	15	27.08	10.40	38.41	14	33.25	16.01	48.16

TABLE 8-continued

Buprenorphine In Vivo Absorption Data for Test Formulation 3

	Suboxone ® sublingual				Test Formulation 3 (pH = 5-5.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
AUC_{inf} (hr * ng/mL)	15	29.58	11.15	37.68	13	38.34	15.38	40.13
$T_{1/2}$ (hr)	15	44.76	20.86	46.60	13	41.71	17.70	42.42

TABLE 9

Naloxone In Vivo Absorption Data for Test Formulation 3

	Suboxone ® sublingual				Test Formulation 3 (pH = 5-5.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
T_{max} (hr)	15	0.90	0.23	25.32	14	0.98	0.62	63.51
C_{max} (pg/mL)	15	94.6	39.1	41.33	14	173	84.5	48.79
AUC_{last} (hr * pg/mL)	15	297.1	120.7	40.62	14	455.2	195.5	42.94
AUC_{inf} (hr * pg/mL)	15	306.1	122.6	40.06	13	474.4	203.1	42.81
$T_{1/2}$ (hr)	15	6.62	2.60	39.26	13	9.45	6.90	73.00

As can be seen, the in vivo data indicated that the absorption of buprenorphine increased as the local pH level decreased. It appeared that by decreasing the local pH from 6.5 to 5.5, the absorption of buprenorphine was being moved to a level further away from that of the one dose tablet. In addition, the naloxone values did not provide a bioequivalent result as the one dose tablet. Thus, it was determined that the film having a local pH of 5.5 did not provide a bioequivalent result as that of the Suboxone® tablet for both buprenorphine and naloxone.

It was noted that by reducing the local pH of the film to a level of 5.5, there would be provided an increased level of absorption of buprenorphine. Thus, it may be desirable to buffer a film composition incorporating buprenorphine itself to a level of about 5.5 to provide an increased absorption.

Example 8

Analysis of In Vivo Absorption of Film Having a pH of 3-3.5

Having determined the absorption of buprenorphine and naloxone in films having a local pH of 6.5 and 5.5, a film dosage composition of film having a local pH of about 3-3.5 was analyzed. It was assumed that the absorption of buprenorphine would continue to be increased as it had demonstrated at a local pH of 5.5. Thus, it was assumed that at a local pH of 3.5, the film would not be bioequivalent to that of the tablet.

Specifically, Test Formulation 2, as prepared in Example 5, was analyzed in vivo to determine the absorption of buprenorphine and of naloxone. The comparative films were compared to the absorption of buprenorphine and of naloxone provided by the Suboxone® one dose tablet. The test film was compared to determine whether it provided a bioequivalent effect as the tablet product.

The results for Test Formulation 2, which had a local pH of about 3-3.5, as compared to the Suboxone® tablet, are set forth in Tables 10 and 11 below.

TABLE 10

Buprenorphine In Vivo Absorption Data for Test Formulation 2

	Suboxone ® sublingual				Test Formulation 2 (pH = 3-3.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
T_{max} (hr)	15	1.60	0.47	29.41	14	1.68	0.58	34.68
C_{max} (ng/mL)	15	2.27	0.562	24.77	14	2.68	0.910	33.99
AUC_{last} (hr * ng/mL)	15	27.08	10.40	38.41	14	29.73	12.05	40.54
AUC_{inf} (hr * ng/mL)	15	29.58	11.15	37.68	14	31.45	12.98	41.26
$T_{1/2}$ (hr)	15	44.76	20.86	46.60	14	30.03	13.95	46.46

TABLE 11

Naloxone In Vivo Absorption Data for Test Formulation 2

	Suboxone ® sublingual				Test Formulation 2 (pH = 3-3.5)			
Parameter	n	Mean	SD	CV %	n	Mean	SD	CV %
T_{max} (hr)	15	0.90	0.23	25.32	14	0.84	0.19	22.19
C_{max} (pg/mL)	15	94.6	39.1	41.33	14	130	72.9	56.04
AUC_{last} (hr * pg/mL)	15	297.1	120.7	40.62	14	362.2	155.9	43.03
AUC_{inf} (hr * pg/mL)	15	306.1	122.6	40.06	12	350.4	142.3	40.61
$T_{1/2}$ (hr)	15	6.62	2.60	39.26	12	8.07	4.75	58.84

As can be seen, the in vivo data indicated that the absorption of buprenorphine was substantially bioequivalent to that of the one dose tablet when the film composition local pH was lowered to about 3-3.5. This result was surprising as it did not appear to follow the pH partition theory. Further, at a local pH of about 3-3.5, it was seen that the absorption of naloxone was substantially bioequivalent to that of the one dose tablet.

Thus, it was determined that the film product including buprenorphine and naloxone at a local pH of 3-3.5 was substantially bioequivalent to that of the Suboxone® one dose tablet.

Example 9

Normalized Values for Naloxone in Films and Tablets

Various film compositions including buprenorphine and naloxone in 8/2 mg and 2/0.5 mg dosages, and having different local pH values from 6.5 to 3.5, were prepared and analyzed. The data was normalized and compared to the one dose tablet. The results are set forth in Table 12 below.

TABLE 12

Normalized Values for Naloxone Film Compared to Tablet

pH	Dose (mg) Buprenorphine/Naloxone	AUC (Normalized)	Cmax	Mg Citric Acid	Ratio Citric Acid(mg)/ Naloxone (mg)
6.5	8/2	3.02	4.33	1.34	0.67
5.5	8/2	1.55	1.83	1.34	0.67
3.5	8/2	1.14	1.37	1.34	0.67
3.5	2/0.5	0.98	0.90	1.34	2.68
5.5	2/0.5	1.41	1.41	1.34	2.68

The data indicates that not only is the local pH of significant importance, but the amount of buffer present in the formula is also important. The improvement from the 8/2 dose to the 2/0.5 dose (at a local pH of 3.5) demonstrates this importance. The 8/2 dose has a ratio of buffer/naloxone of 0.67, and this dose provided borderline acceptable bioequivalent results. In contrast, the 2/0.5 dose has a ratio of buffer/naloxone of 2.68, and provides a more bioequivalent absorption value than the 8/2 dose.

In fact, the data shows that the 2/0.5 dose at a local pH of 3.5 had an even lower buccal absorption than the one dose tablet, as seen from the normalized values for the AUC and Cmax. This demonstrates that even less absorption of the naloxone occurs for the film formulation at a local pH of 3.5 than the tablet formulation. Given the goal of reducing the absorption of naloxone, it appears that the film product buffered at a local pH of 3.5 with a buffer ratio of buffer/Naloxone of 2.68 provides even better results than the Suboxone® tablet formulation.

What is claimed is:

1. A film dosage composition comprising:

a. A polymeric carrier matrix;

b. A therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof;

c. A therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof; and

d. A buffer in an amount to provide a local pH for said composition of a value sufficient to optimize absorption of said buprenorphine, wherein said local pH is from about 3 to about 3.5 in the presence of saliva.

2. The composition of claim **1**, wherein said film dosage composition provides a bioequivalent absorption of buprenorphine to that of a tablet having an equivalent amount of buprenorphine or a pharmaceutically acceptable salt thereof.

3. The composition of claim **1**, wherein said polymeric carrier matrix comprises at least one polymer in an amount of at least 25% by weight of said composition.

4. The composition of claim **1**, wherein said buffer is present in an amount of from about 2:1 to about 1:5 by weight of buffer to buprenorphine.

5. The composition of claim **1**, wherein said polymeric carrier matrix comprises at least one self-supporting film forming polymer.

6. The film dosage composition of claim **1**, wherein said buprenorphine is present in an amount of from about 2 mg to about 16 mg per dosage.

7. The film dosage composition of claim **1**, wherein said buffer comprises sodium citrate, citric acid, and combinations thereof.

8. The film dosage composition of claim **1**, wherein said buffer comprises acetic acid, sodium acetate, and combinations thereof.

9. A method of treating narcotic dependence of a user, comprising the steps of:

a. providing a composition comprising:

i. A polymeric carrier matrix;

ii. A therapeutically effective amount of buprenorphine or a pharmaceutically acceptable salt thereof;

iii. A therapeutically effective amount of naloxone or a pharmaceutically acceptable salt thereof; and

iv. A buffer in an amount to provide a local pH of about 3 to about 3.5 for said composition of a value sufficient to optimize absorption of said buprenorphine and also sufficient to inhibit absorption of said naloxone; and

b. administering said composition to the oral cavity of a user.

10. The composition of claim **9**, wherein said method provides a bioequivalent absorption of buprenorphine to that of a tablet having an equivalent amount of buprenorphine or a pharmaceutically acceptable salt thereof.

11. The method of claim **9**, wherein said film dosage composition is administered to the user through buccal administration, sublingual administration, and combinations thereof.

12. The method of claim **9**, wherein said film dosage composition remains in the oral cavity of the user for a period of at least 1 minute.

13. The method of claim **9**, wherein said film dosage composition remains in the oral cavity of the user for a period of between about 1 and 1.5 minutes.

14. The method of claim **9**, wherein said film dosage composition remains in the oral cavity of the user for a period of up to 3 minutes.

15. An orally dissolving film formulation comprising buprenorphine and naloxone, wherein said formulation provides an in vivo plasma profile having a Cmax of between about 0.624 ng/ml and about 5.638 ng/ml for buprenorphine and an in vivo plasma profile having a Cmax of between about 41.04 pg/ml to about 323.75 pg/ml for naloxone.

16. The formulation of claim **15**, wherein said formulation provides a mean AUC of between about 5.431 hr·ng/ml to about 56.238 hr·ng/ml for buprenorphine.

17. The formulation of claim **15**, wherein said formulation provides a mean AUC of between about 102.88 hr·pg/ml to about 812.00 hr·pg/ml for naloxone.

18. The formulation of claim **15**, wherein said formulation comprises about 2 to about 16 mg of buprenorphine or a salt thereof.

19. The formulation of claim **15**, wherein said formulation comprises about 0.5 to about 4 mg of naloxone or a salt thereof.

* * * * *

EXHIBIT B

US008017150B2

(12) **United States Patent**
Yang et al.

(10) **Patent No.: US 8,017,150 B2**
(45) **Date of Patent: Sep. 13, 2011**

(54) **POLYETHYLENE OXIDE-BASED FILMS AND DRUG DELIVERY SYSTEMS MADE THEREFROM**

(75) Inventors: **Robert K. Yang**, Flushing, NY (US); **Richard C. Fuisz**, McLean, VA (US); **Garry L. Myers**, Kingsport, TN (US); **Joseph M. Fuisz**, Washington, DC (US)

(73) Assignee: **MonoSol Rx, LLC**, Portage, IN (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 364 days.

(21) Appl. No.: **12/107,389**

(22) Filed: **Apr. 22, 2008**

(65) **Prior Publication Data**

US 2008/0260809 A1 Oct. 23, 2008

Related U.S. Application Data

(60) Division of application No. 10/856,176, filed on May 28, 2004, now Pat. No. 7,666,337, which is a continuation-in-part of application No. PCT/US02/032575, filed on Oct. 11, 2002, and a continuation-in-part of application No. PCT/US02/32594, filed on Oct. 11, 2002, and a continuation-in-part of application No. PCT/US02/32542, filed on Oct. 11, 2002.

(60) Provisional application No. 60/473,902, filed on May 28, 2003, provisional application No. 60/414,276, filed on Sep. 27, 2002, provisional application No. 60/371,940, filed on Apr. 11, 2002.

(51) **Int. Cl.**
A61K 9/14 (2006.01)

(52) **U.S. Cl.** **424/484**; 424/486; 424/488; 424/434; 424/435

(58) **Field of Classification Search** 424/434, 424/435, 436, 443, 484
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

307,537	A	11/1884	Foulks
688,446	A	12/1901	Stempel
2,142,537	A	1/1939	Tisza
2,277,038	A	3/1942	Curtis
2,352,691	A	7/1944	Curtis
2,501,544	A	3/1950	Shrontz
2,980,554	A	4/1961	Gentile et al.
3,249,109	A	5/1966	Maeth et al.
3,444,858	A	5/1969	Russell
3,536,809	A	10/1970	Applezweig
3,551,556	A	12/1970	Kliment et al.
3,598,122	A	8/1971	Zaffaroni
3,632,740	A	1/1972	Robinson et al.
3,640,741	A	2/1972	Etes
3,641,237	A	2/1972	Gould et al.
3,731,683	A	5/1973	Zaffaroni
3,753,732	A	8/1973	Boroshok
3,814,095	A	6/1974	Lubens
3,892,905	A	7/1975	Albert
3,911,099	A	10/1975	DeFoney et al.
3,972,995	A	8/1976	Tsuk et al.
3,996,934	A	12/1976	Zaffaroni
3,998,215	A	12/1976	Anderson et al.
4,029,757	A	6/1977	Mlodozeniec et al.
4,029,758	A	6/1977	Mlodozeniec et al.
4,031,200	A	6/1977	Reif
4,123,592	A	10/1978	Rainer et al.
4,128,445	A	12/1978	Sturzenegger et al.
4,136,145	A	1/1979	Fuchs et al.
4,136,162	A	1/1979	Fuchs et al.
4,139,627	A	2/1979	Lane et al.
4,226,848	A	10/1980	Nagai et al.
4,251,400	A	2/1981	Columbus
4,292,299	A	9/1981	Suzuki et al.
4,294,820	A	10/1981	Keith et al.
4,302,465	A	11/1981	Ekenstam et al.
4,307,075	A	12/1981	Martin
4,325,855	A	4/1982	Dickmann et al.
4,373,036	A	2/1983	Chang et al.
4,406,708	A	9/1983	Hesselgren
4,432,975	A	2/1984	Libby
4,438,258	A	3/1984	Graham
4,460,562	A	7/1984	Keith et al.
4,466,973	A	8/1984	Rennie
4,503,070	A	3/1985	Eby, III
4,515,162	A	5/1985	Yamamoto et al.
4,517,173	A	5/1985	Kizawa et al.
4,529,601	A	7/1985	Broberg et al.
4,529,748	A	7/1985	Wienecke
4,562,020	A	12/1985	Hijiya et al.
4,569,837	A	2/1986	Suzuki et al.
4,593,053	A	6/1986	Jevne et al.
4,608,249	A	8/1986	Otsuka et al.
4,615,697	A	10/1986	Robinson
4,623,394	A	11/1986	Nakamura et al.

(Continued)

FOREIGN PATENT DOCUMENTS

DE	2432925	C3	1/1976

(Continued)

OTHER PUBLICATIONS

Flick, E., Water-Soluble Resins—An Industrial Guide, 1991, (2nd Ed.) William Andrew Publishing/Noyes, pp. 389-391.*

(Continued)

Primary Examiner — Gina C Yu
(74) *Attorney, Agent, or Firm* — Hoffmann & Baron, LLP

(57) **ABSTRACT**

The invention relates to the film products and methods of their preparation that demonstrate a non-self-aggregating uniform heterogeneity. Desirably, the films disintegrate in water and may be formed by a controlled drying process, or other process that maintains the required uniformity of the film. The films contain a polymer component, which includes polyethylene oxide optionally blended with hydrophilic cellulosic polymers. Desirably, the films also contain a pharmaceutical and/or cosmetic active agent with no more than a 10% variance of the active agent pharmaceutical and/or cosmetic active agent per unit area of the film.

18 Claims, 34 Drawing Sheets

U.S. PATENT DOCUMENTS

4,652,060	A		3/1987	Miyake
4,659,714	A		4/1987	Watt-Smith
4,675,009	A		6/1987	Hymes et al.
4,695,465	A		9/1987	Kigasawa et al.
4,704,119	A		11/1987	Shaw et al.
4,713,239	A		12/1987	Babaian et al.
4,713,243	A		12/1987	Schiraldi et al.
4,722,761	A		2/1988	Cartmell et al.
4,740,365	A		4/1988	Yukimatsu et al.
4,748,022	A		5/1988	Busciglio
4,765,983	A		8/1988	Takayanagi et al.
4,772,470	A		9/1988	Inoue et al.
4,777,046	A		10/1988	Iwakura et al.
4,789,667	A		12/1988	Makino et al.
4,849,246	A		7/1989	Schmidt
4,860,754	A		8/1989	Sharik et al.
RE33,093	E		10/1989	Schiraldi et al.
4,876,092	A		10/1989	Mizobuchi et al.
4,876,970	A		10/1989	Bolduc
4,880,416	A		11/1989	Horiuchi et al.
4,888,354	A		12/1989	Chang et al.
4,894,232	A		1/1990	Reul et al.
4,900,552	A		2/1990	Sanvordeker et al.
4,900,554	A		2/1990	Yanagibashi et al.
4,900,556	A		2/1990	Wheatley et al.
4,910,247	A		3/1990	Haldar et al.
4,915,950	A		4/1990	Miranda et al.
4,925,670	A		5/1990	Schmidt
4,927,634	A		5/1990	Sorrentino et al.
4,927,636	A		5/1990	Hijiya et al.
4,937,078	A		6/1990	Mezei et al.
4,940,587	A		7/1990	Jenkins et al.
4,948,580	A		8/1990	Browning
4,958,580	A		9/1990	Asaba et al.
4,978,531	A		12/1990	Yamazaki et al.
4,981,693	A		1/1991	Higashi et al.
4,981,875	A		1/1991	Leusner et al.
5,023,082	A		6/1991	Friedman et al.
5,024,701	A		6/1991	Desmarais
5,028,632	A		7/1991	Fuisz
5,047,244	A		9/1991	Sanvordeker et al.
5,064,717	A		11/1991	Suzuki et al.
5,089,307	A		2/1992	Ninomiya et al.
5,158,825	A		10/1992	Altwirth
5,166,233	A		11/1992	Kuroya et al.
5,186,938	A		2/1993	Sablotsky et al.
5,229,164	A		7/1993	Pins et al.
5,234,957	A		8/1993	Mantelle
5,271,940	A		12/1993	Cleary et al.
5,272,191	A		12/1993	Ibrahim et al.
5,346,701	A		9/1994	Heiber et al.
5,393,528	A		2/1995	Staab
5,411,945	A		5/1995	Ozaki et al.
5,413,792	A		5/1995	Ninomiya et al.
5,433,960	A		7/1995	Meyers
5,455,043	A		10/1995	Fischel-Ghodsian
5,462,749	A		10/1995	Rencher
5,472,704	A		12/1995	Santus et al.
5,518,902	A		5/1996	Ozaki et al.
5,567,431	A		10/1996	Vert et al.
5,620,757	A		4/1997	Ninomiya et al.
5,629,003	A		5/1997	Horstmann et al.
5,656,296	A	*	8/1997	Khan et al. 424/473
5,700,478	A		12/1997	Biegajski et al.
5,700,479	A		12/1997	Lundgren
5,766,620	A		6/1998	Heiber et al.
5,800,832	A		9/1998	Tapolsky et al.
5,948,430	A		9/1999	Zerbe et al.
6,072,100	A		6/2000	Mooney et al.
6,153,210	A		11/2000	Roberts et al.
6,177,096	B1		1/2001	Zerbe et al.
6,231,957	B1		5/2001	Zerbe et al.
6,264,981	B1		7/2001	Zhang et al.
6,284,264	B1		9/2001	Zerbe et al.
6,375,963	B1		4/2002	Repka et al.
6,488,963	B1		12/2002	McGinity et al.
6,800,329	B2		10/2004	Horstmann et al.
7,579,019	B2		8/2009	Tapolsky et al.
2001/0006677	A1		7/2001	McGinity et al.
2001/0022964	A1		9/2001	Leung et al.
2001/0046511	A1		11/2001	Zerbe et al.
2003/0069263	A1		4/2003	Breder et al.
2004/0191302	A1		9/2004	Davidson
2005/0048102	A1		3/2005	Tapolsky et al.
2005/0118217	A1		6/2005	Barnhart et al.
2007/0087036	A1		4/2007	Durschlag et al.
2007/0148097	A1		6/2007	Finn et al.
2008/0254105	A1		10/2008	Tapolsky et al.

FOREIGN PATENT DOCUMENTS

DE	2449865	B2	4/1976
DE	3630603	C2	3/1988
DE	19646392	A1	5/1998
EP	0200508	B1	12/1986
EP	0219762	A1	4/1987
EP	0241178	B1	10/1987
EP	0250187	B1	12/1987
EP	0259749	B1	3/1988
EP	0273069	B1	7/1988
EP	0381194	A2	8/1990
EP	0452446	B1	10/1991
EP	0514691	B1	11/1992
EP	0598606	A1	5/1994
EP	1110546	A1	6/2001
WO	9105540		5/1991
WO	9215289		9/1992
WO	9505416		2/1995
WO	9518046		7/1995
WO	9731621		9/1997
WO	0018365		4/2000
WO	0042992		7/2000
WO	0170194	A1	9/2001
WO	0191721	A2	12/2001
WO	03030882	A1	4/2003
WO	03030883	A1	4/2003
WO	2005102287		11/2005
WO	2008011194	A2	1/2008

OTHER PUBLICATIONS

XP-002298105; Polyethylenglykole; Internet: www.roempp.com; Sep. 20, 2004.

Repka et al.; Influence of Vitamin E. TPGS on the properties of hydrophilic films produced by hot-melt extrusion; International Journal of Pharmaceutics; vol. 202, pp. 63-70; 2000.

Repka et al.; Bioadhesive properties of hydroxypropylcellulose topical films produced by hot-melt extrusion; Journal of Controlled Release; vol. 70; pp. 341-351; 2001.

International Search Report for PCT/US2004/017076.

* cited by examiner




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30





FIG. 31



Normalized to GAPDH, most of the ISGs are induced.

FIG. 32



PROBE TEMP (C)
OVEN TEMP (C)
90
80
70
60
50
40
30
20
10
0
1
2
3
4
5
6
7
8
9
10
11
12

FIG. 33



PROBE TEMP (C)
OVEN TEMP (C)
90
80
70
60
50
40
30
20
10
0
1
2
3
4
5
6
7
8
9
10
11
12

FIG. 34




FIG. 35




FIG. 36




FIG. 37

Ex.	Polymer Component Reference	% Solids of solution	Viscosity (cp) at 5 rpm	% moisture	Film thickness (mils)	Film strength	Tear Resistance	Tendency to go to roof of mouth	180° bend test	Film molding	Dis-solution (sec)	Rating of dissolution in mouth	Time in oven (min)
EI	PEO/PVP (60/40)	45.0	14800	2.21	3.8	Adequate	Excellent	Low	Passed	No	3	Fast to Moderate	9
EJ	PEO/PVP (40/60)	50.0	6600	2.86	4	Weak	Low to moderate	High	Passed	No	3	Fast	8
EK	PEO/Starch (80/20)	40.0	3440	2.27	4.5	Adequate to good	Excellent	High	Passed	No	3	Fast to Moderate	8
EL	PEO/CMC (80/20)	37.5	121,200	1.96	4.1	Good	Excellent	High	Passed	No	5	Slow	9
EM	PEO/CMC (60/40)	30.0	82,000	4.21	3.45	Weak	Good	High	Passed	No	3	Slow to Moderate	9
EN	PEO/CMC (40/60)	30.0	185,000	3.07	3.5	Adequate	Very low	High	Failed	No	4	Slow	9
EO	PEO/HPC (80/20)	37.5	21,200	1.65	4	Good	Excellent	High	Passed	No	4	Fast	8
EP	PEO/HPC (60/40)	37.5	17,000	2.84	3.8	Adequate	Excellent	High	Passed	No	4	Fast	9
EQ	PEO/HPC (40/60)	42.5	43,400	2.83	4.5	Poor to adequate	Poor to good	High	Passed	No	7	Fast to Moderate	7
ER	PEO/HPC (20/80)	42.5	46,400	2.33	4.4	Adequate to good	Poor	Low	Passed	No	14-15	Slow	9
ES	PEO/HPMC (80/20)	37.5	29,000	2.14	4.4	Adequate	Good	High	Passed	Yes	4	Fast to Moderate	8
ET	PEO/HPMC (60/40)	37.5	47,000	2.37	3.9	Poor to adequate	Slight	High	Passed	Yes	3	Fast to Moderate	9
EU	PEO/HPMC (40/60)	35.0	54,800	3.55	4.5	Adequate to good	Low	Low	Passed	Yes	8	Slow	8
EV	PEO/HPMC (20/80)	35.0	96,600	4.43	4.5	Good	Low	Low	Passed	No	22	Slow	10
EW	PEO/PVA (80/20)	37.5	41,600	2.92	9	Weak	Moderate	High	Passed	No	3	Moderate	10

FIG. 38