EXHIBIT

I

US006627210B2

(12) **United States Patent**
Olejnik et al.

(10) **Patent No.: US 6,627,210 B2**
(45) **Date of Patent: *Sep. 30, 2003**

(54) **COMPOSITIONS CONTAINING α-2-ADRENERGIC AGONIST COMPONENTS**

(75) Inventors: **Orest Olejnik**, Coto de Coza, CA (US); **Edward D. S. Kerslake**, Charlestown, MA (US)

(73) Assignee: **Allergan, Inc.**, Irvine, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 8 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **09/904,018**

(22) Filed: **Jul. 10, 2001**

(65) **Prior Publication Data**

US 2002/0032201 A1 Mar. 14, 2002

Related U.S. Application Data

(60) Provisional application No. 60/218,200, filed on Jul. 14, 2000.

(51) **Int. Cl.**[7] **A61K 33/14**; A61K 9/00; A61K 9/08

(52) **U.S. Cl.** **424/427**; 424/400; 424/401; 424/466; 424/422; 514/772.4; 514/772.6

(58) **Field of Search** 424/427, 400, 424/422, 661, 407; 514/772.4, 772.6

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,278,447	A		10/1966	McNicholas
3,890,319	A		6/1975	Danielwicz et al.
4,530,920	A		7/1985	Nestor et al.
4,806,556	A		2/1989	Portoghese
5,021,416	A		6/1991	Gluchowski
5,202,128	A		4/1993	Morella et al.
5,215,991	A	*	6/1993	Burke 514/255
5,352,796	A		10/1994	Hoeger et al.
5,459,133	A	*	10/1995	Neufeld 514/211
5,703,077	A		12/1997	Burke et al.
5,719,197	A		2/1998	Kanios et al.
5,725,887	A		3/1998	Martin et al.
5,814,638	A		9/1998	Lee et al.
5,834,502	A		11/1998	Cheng et al.
5,994,110	A		11/1999	Mosbach et al.
2002/0010202	A1	*	1/2002	Garst 514/392
2002/0071874	A1	*	6/2002	Olejnik et al. 424/661

FOREIGN PATENT DOCUMENTS

CA	2048315	2/1992
EP	0609961	8/1994
WO	94/16685	8/1994
WO	98/47878	10/1998
WO	99/43299	9/1999
WO	99/51273	10/1999
WO	0012137	3/2000
WO	0019981	4/2000

OTHER PUBLICATIONS

Remington's Pharmaceutical Sciences, Eighteenth Edition, 1990, pp. 1304–1305.*
U.S. patent application Ser. No. 09/903,962, Olejnik et al., filed Jul. 10, 2001.

* cited by examiner

Primary Examiner—Thurman K. Page
Assistant Examiner—Rachel M. Bennett
(74) *Attorney, Agent, or Firm*—Carlos A. Fisher; Martin A. Voet; Robert J. Baran

(57) **ABSTRACT**

Compositions useful for improving effectiveness of alpha-2-adrenergic agonist components include carrier components, alpha-2-adrenergic agonist components, solubility enhancing components which aid in solubilizing the alpha-2-adrenergic agonist components. In one embodiment, the alpha-2-adrenergic agonist components include alpha-2-adrenergic agonists. In another embodiment, the solubility enhancing components include carboxymethylcellulose.

34 Claims, 1 Drawing Sheet




SOLUBILITY OF BRIMONIDINE TARTRATE (ppm)
4000
3500
3000
2500
2000
1500
1000
500
0
7
7.5
8
8.5
9
9.5
10
10.5
pH
0% CMC
0.058% CMC
0.17% CMC
0.5% CMC
1.5% CMC

COMPOSITIONS CONTAINING α-2-ADRENERGIC AGONIST COMPONENTS

CROSS REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application 60/218,200 filed Jul. 14, 2000.

BACKGROUND OF THE INVENTION

The present invention relates to compositions containing alpha-2-adrenergic agonist components. More particularly, the invention relates to such compositions in which the alpha-2-adrenergic agonist components have enhanced solubility at the therapeutically effective concentrations.

Alpha-2-adrenergic agonist components include chemical entities, such as compounds, ions, complexes and the like, which are effective to act on or bind to Alpha-2-adrenergic receptors and provide a therapeutic effect. Alpha-2-adrenergic agonist components means the agonists themselves and any and all precursors thereof, metabolites thereof and combinations thereof. One of the continuing challenges of formulating compositions having alpha-2-adrenergic agonist components is to render such components more effective. For example, alpha-2-adrenergic agonist components in liquid compositions often benefit from being soluble in the liquid carriers of such compositions. Such solubility promotes uniform and accurate administration.

Additionally, the dispensed or administered alpha-2-adrenergic agonist components should advantageously be soluble in biological systems or environments, for example, for effective or enhanced in vivo diffusion through cell membranes or lipid bilayers. Some alpha-2-adrenergic agonist components with higher pKa's, for example, greater than about 7, tend to diffuse very well through lipid membranes at pH valves near their pka, because in such circumstances they are predominantly unionized in neutral to alkaline biological environments. However, some of these alpha-2-adrenergic agonist components become insoluble at neutral to alkaline biological pH's. Such insolubility may decrease membrane diffusion capabilities, rendering the alpha-2-adrenergic agonist components less effective and/or their therapeutic effects more variable at a given dosage. Furthermore, solubilized alpha-2-adrenergic agonist components provide other benefits, for example, reduced irritation to tissues that interact with alpha-2-adrenergic agonist components.

There continues to be a need for new compositions containing alpha-2-adrenergic agonist components.

SUMMARY OF THE INVENTION

New alpha-2-adrenergic agonist component-containing compositions have been discovered. The present compositions contain certain materials which are effective in at least aiding or assisting in solubilizing the alpha-2-adrenergic agonist components in the compositions, and preferably in environments to which the compositions are administered or introduced, for example, biological environments, such as the human eye. Preferably, solubilization of the alpha-2-adrenergic agonist components in accordance with the present invention facilitates transport of such components across lipid membranes. Also, preferably such solubilization allows the provision of more reliable and reproducible dosage forms of the drug. In addition, alpha-2-adrenergic agonist component-containing compositions have been discovered which include preservatives which provide substantial advantages, for example, reduced adverse interactions with the alpha-2-adrenergic agonist components and/or with the patients to whom the compositions are administered, while maintaining preservative effectiveness.

The present compositions preferably enhance the effectiveness of alpha-2-adrenergic agonist components by increasing the apparent water solubility of the alpha-2-adrenergic agonist components, preferably at pH's higher than neutral. The present compositions include, in addition to the adrenergic agonist components, solubility enhancing components (SECs) in amounts effective to enhance the solubility of the alpha-2-adrenergic agonist components. Preferably, the alpha-2-adrenergic agonist components are more soluble in the present compositions having, for example, pH's of about 7 or greater, relative to similar compositions without the SECs. In another embodiment, the alpha-2-adrenergic agonist components of the present compositions are more soluble in neutral, preferably alkaline, biological environments into which the compositions are administered relative to alpha-2-adrenergic agonist components in similar compositions without the SECs.

In one embodiment, the alpha-2-adrenergic agonist components include imino-imidazolines, imidazolines, imidazoles, azepines, thiazines, oxazolines, guanidines, catecholamines, biologically compatible salts and esters and mixtures thereof. Preferably, the alpha-2-adrenergic agonist components include quinoxaline components. Quinoxaline components include quinoxaline, biologically compatible salts thereof, esters thereof, other derivatives thereof and the like, and mixtures thereof. Non-limiting examples of quinoxaline derivatives include (2-imidozolin-2-ylamino) quinoxaline, 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, and biologically compatible salts thereof and esters thereof, preferably the tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, and the like and mixtures thereof. Hereinafter, the tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline is referred to as "Brimonidine tartrate."

In a preferred embodiment, the alpha-2-adrenergic agonist components, such as those listed above, are specific for the alpha-2A-adrenergic receptors, alpha-2B-adrenergic receptors and/or alpha-2D-adrenergic receptors.

In one embodiment, the alpha-2-adrenergic agonist components are unionized in the compositions. Preferably, the alpha-2-adrenergic agonist components are also unionized in the biological environment into which the compositions are administered.

In a useful embodiment, the SEC includes a polyanionic component. As used herein, the term "polyanionic component" refers to a chemical entity, for example, an ionically charged species, such as an ionically charged polymeric material, which includes more than one discrete anionic charge, that is multiple discrete anionic charges. Preferably, the polyanionic component is selected from polymeric materials having multiple anionic charges, and mixtures thereof.

Particularly useful polyanionic components are selected from anionic polymers derived from acrylic acid (meaning to include polymers from acrylic acid, acrylates and the like and mixtures thereof), anionic polymers derived from methacrylic acid (meaning to include polymers from methacrylic acid, methacrylates, and the like and mixtures thereof), anionic polymers derived from alginic acid (meaning to include alginic acid, alginates, and the like and mixtures thereof), anionic polymers of amino acids (meaning to include polymers of amino acids, amino acid salts, and the like and mixtures thereof), and the like, and mixtures

thereof. Very useful polyanionic components are those selected from anionic cellulose derivatives and mixtures thereof, especially carboxymethylcelluloses.

The polyanionic component preferably is sufficiently anionic to interact with or otherwise affect, in particular increase, the solubility of the alpha-2-adrenergic components. This interaction preferably is sufficient to render the alpha-2-adrenergic components substantially completely soluble at therapeutically effective concentrations. The amount of SEC in the composition preferably is in the range of about 0.1% (w/v) to about 30% (w/v), more preferably about 0.2% (w/v) to about 10% (w/v), and even more preferably about 0.2% (w/v) to about 0.6% (w/v).

The compositions include carrier components, for example, aqueous liquid carrier components. In one embodiment, the compositions have pH's of about 7 or greater, preferably about 7 to about 9, and are ophthalmically acceptable.

In a preferred embodiment, a composition is provided which includes an alpha-2-adrenergic agonist component in an amount effective to provide at least one therapeutic benefit to a patient to whom the composition is administered, an anionic cellulose derivative in an amount effective to increase the solubility of the alpha-2-adrenergic agonist component and an aqueous liquid carrier component. The alpha-2-adrenergic agonist component preferably comprises a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline. The anionic cellulose derivative preferably comprises a carboxymethylcellulose. The concentration of the anionic cellulose derivative in the composition should be about 0.2% (w/v) to about 0.6% (w/v).

In a preferred embodiment, the present compositions are ophthalmically acceptable, e.g. the compositions do not have deleterious or toxic properties which could harm the eye of the human or animal to whom the compositions are administered.

In one broad aspect of the invention, complexes are formed in the compositions. In one embodiment, the complexes include monomer units derived from at least one quinoxaline component. In a preferred embodiment, the complexes of the present invention are dimers. In a particularly preferred embodiment, the complexes are complexes, especially dimers, of Bromodidine tartrate.

In another broad aspect of the present invention, compositions are provided which comprise an alpha-2-adrenergic agonist component and a preservative component in an effective amount to at least aid in preserving the compositions. Preferably, the preservative components include oxychloro components, such as compounds, ions, complexes and the like which are biologically acceptable, chemically stable and do not substantially or significantly detrimentally affect the an alpha-2-adrenergic agonist component in the compositions or the patients to whom the compositions are administered. Such compositions preferably are substantially free of cyclodextrins in the compositions or the patients to whom the compositions are administered.

Any feature or combination of features described herein are included within the scope of the present invention provided that the features included in any such combination are not mutually inconsistent as will be apparent from the context, this specification, and the knowledge of one of ordinary skill in the art.

Additional advantages and aspects of the present invention are apparent in the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 is a graph of soluble Brimonidine tartrate verses pH at various carboxymethylcellulose concentrations.

DETAILED DESCRIPTION OF THE INVENTION

Compositions comprising alpha-2-adrenergic agonist components and SECs are provided. The alpha-2-adrenergic agonist components in the present compositions are made more soluble and may be more effectively utilized as therapeutic agents. The SECs employed in the present compositions may be effective in the solubilization of ionized alpha-2-adrenergic agonist components, unionized alpha-2-adrenergic agonist components or both. The present compositions include liquid carrier components and have the characteristics of liquid, for example, aqueous liquid, solutions.

Preferably, the alpha-2-adrenergic agonist components have increased solubility in the present compositions at pH's greater than 7, as compared to identical alpha-2-adrenergic agonist components, at comparable concentrations, in similar compositions without the SECs. More preferably, the alpha-2-adrenergic agonist components have increased solubility in the present compositions at pH's in the range of about 7 to about 10 and, as compared to identical alpha-2-adrenergic agonist components in similar compositions, at comparable concentrations, without the SECs.

Without wishing to be limited by any theory or mechanism of operation, it is believed that solubilized alpha-2-adrenergic agonist components are better able to cross the lipid membranes relative to unsolubilized alpha-2-adrenergic agonist components. It is further believed that the solubilized alpha-2-adrenergic agonist components are physically smaller and are therefore more able to physically permeate or diffuse through the lipid membranes.

In one embodiment, the SECs of this invention are capable of solubilizing the alpha-2-adrenergic agonist components in the biological environments into which they are introduced at therapeutically effective concentrations. Preferably, the biological environments into which the present compositions are introduced have pH's ranging from about 7 to about 9. For example, a composition comprising a SEC and an alpha-2-adrenergic agonist component may be administered to the cornea of an eye, which has a pH of about 7, wherein the alpha-2-adrenergic agonist component is substantially solubilized at the administered area. Furthermore, in one embodiment, the alpha-2-adrenergic agonist components solubilized by SECs at the administered area diffuse through biological lipid membranes more readily than alpha-2-adrenergic agonist components which are not solubilized by SECs. The solubilization of alpha-2-adrenergic agonist components preferably reduces irritation to sensitive tissues in contact or interacting with the alpha-2-adrenergic agonist components.

The presently useful alpha-2-adrenergic agonist components preferably are chosen to benefit from the presence of the SECs. In general, the alpha-2-adrenergic agonist components are provided with increased apparent solubility, preferably increased apparent water solubility, by the presence of the SECs.

Examples of alpha-2-adrenergic agonist components include molecules containing amines. Preferably, the alpha-2-adrenergic agonist components are amine-containing molecules with pKa's of greater than about 7, more preferably about 7 to about 9.

Alpha-2-adrenergic agonist components include alpha-2-adrenergic agonists. As used herein, the term alpha-2 adrenergic agonist includes chemical entities, such as compounds, ions, complexes and the like, that produce a net sympatholytic response, resulting in increased

accommodation, for example, by binding to presynaptic alpha-2 receptors on sympathetic postganglionic nerve endings or for example, to postsynaptic alpha-2 receptors on smooth muscle cells. A sympatholytic response is characterized by the inhibition, diminishment, or prevention of the effects of impulses conveyed by the sympathetic nervous system. The alpha-2 adrenergic agonists of the invention bind to the alpha-2 adrenergic receptors presynaptically, causing negative feedback to decrease the release of neuronal norepinephrine. Additionally, they also work on alpha-2 adrenergic receptors postsynaptically, inhibiting beta-adrenergic receptor-stimulated formation of cyclic AMP, which contributes to the relaxation of the ciliary muscle, in addition to the effects of postsynaptic alpha-2 adrenergic receptors on other intracellular pathways. Activity at either pre- or postsynaptic alpha-2 adrenergic receptors will result in a decreased adrenergic influence. Decreased adrenergic influence results in increased contraction resulting from cholinergic innervations. Alpha-2 adrenergic agonists also include compounds that have neuroprotective activity. For example, 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline is an alpha-2-adrenergic agonist which has a neuroprotective activity through an unknown mechanism.

Without limiting the invention to the specific groups and compounds listed, the following is a list of representative alpha-2 adrenergic agonists useful in this invention: iminoimidazolines, including clonidine, apraclonidine; imidazolines, including naphazoline, xymetazoline, tetrahydrozoline, and tramazoline; imidazoles, including detomidine, medetomidine, and dexmedetomidine; azepines, including B-HT 920 (6-allyl-2-amino-5,6,7,8 tetrahydro-4H-thiazolo[4,5-d]-azepine and B-HT 933; thiazines, including xylazine; oxazolines, including rilmenidine; guanidines, including guanabenz and guanfacine; catecholamines; and the like and derivatives thereof.

Particularly useful alpha-2-adrenergic agonists include quinoxaline components. In one embodiment, the quinoxaline components include quinoxaline, derivatives thereof and mixtures thereof. Preferably, the derivatives of quinoxaline include (2-imidozolin-2-ylamino) quinoxaline. More preferably, the derivatives of quinoxaline include 5-halide-6-(2-imidozolin-2-ylamino) quinoxaline. The "halide" of the 5-halide-6-(2-imidozolin-2-ylamino) quinoxaline may be a fluorine, a chlorine, an iodine, or preferably, a bromine, to form 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline. Even more preferably, the derivatives of quinoxaline to be used in accordance with this invention include a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, or Brimonidine tartrate.

Other useful quinoxaline derivatives are well known. For example, useful derivatives of a quinoxaline include the ones disclose by Burke et al U.S. Pat. No. 5,703,077. See also Danielwicz et al U.S. Pat. No. 3,890,319. Each of the disclosures of Burke et al and Danielwicz et al is incorporated in its entirety by reference herein.

The quinoxaline and derivatives thereof, for example Brimonidine tartrate, are amine-containing and preferably have pKa's of greater than 7, preferably about 7.5 to 9.

Analogs of the foregoing compounds that function as alpha-2 adrenergic agonists also are specifically intended to be embraced by the invention.

Preferably, the alpha-2-adrenergic agonists, for example the ones listed above, are effective toward activating alpha-2A-adrenergic receptors, alpha-2B-adrenergic receptors and alpha-2D-adrenergic receptors.

In one embodiment, the alpha-2-adrenergic agonists, for example Brimonidine tartrate, are substantially unionized in the compositions. In another embodiment, the adrenergic compounds are substantially unionized in the environment to which they are administered, for example the cornea. Without wishing to be limited by any theory or mechanism of action, it is believed that the unionized forms of the adrenergic compounds facilitate their permeation across membrane lipid bilayers.

Any suitable SEC may be employed in accordance with the present invention. In one embodiment, the SECs include pyrrolinidone components. Examples of pyrrolinidone components are polyvinylpyrrolinidones and derivatives thereof. In a preferred embodiment, the SECs include polyanionic components. The useful polyanionic components include, but are not limited to, those materials which are effective in increasing the apparent solubility, preferably water solubility, of poorly soluble alpha-2-adrenergic agonist components and/or enhance the stability of the alpha-2-adrenergic agonist components and/or reduce unwanted side effects of the alpha-2-adrenergic agonist components. Furthermore, the polyanionic component is preferably ophthalmically acceptable at the concentrations used. Additionally, the polyanionic component preferably includes three (3) or more anionic (or negative) charges. In the event that the polyanionic component is a polymeric material, it is preferred that each of the repeating units of the polymeric material include a discrete anionic charge. Particularly useful anionic components are those which are water soluble, for example, soluble at the concentrations used in the presently useful liquid aqueous media, such as a liquid aqueous medium containing the alpha-2-adrenergic components.

The polyanionic component is preferably sufficiently anionic to interact with the alpha-2-adrenergic agonist component. Such interaction is believed to be desirable to solubilize the alpha-2-adrenergic agonist component and/or to maintain such alpha-2-adrenergic agonist component soluble in the carrier component, for example a liquid medium.

Polyanionic components also include one or more polymeric materials having multiple anionic charges. Examples include:

- metal carboxymethylstarchs
- metal carboxymethylhydroxyethylstarchs
- hydrolyzed polyacrylamides and polyacrylonitriles heparin
- homopolymers and copolymers of one or more of:
  - acrylic and methacrylic acids
  - metal acrylates and methacrylates
  - alginic acid
  - metal alginates
  - vinylsulfonic acid
  - metal vinylsulfonate
  - amino acids, such as aspartic acid, glutamic acid and the like
  - metal salts of amino acids
  - p-styrenesulfonic acid
  - metal p-styrenesulfonate
  - 2-methacryloyloxyethylsulfonic acids
  - metal 2-methacryloyloxethylsulfonates
  - 3-methacryloyloxy-2-hydroxypropylsulonic acids
  - metal 3-methacryloyloxy-2-hydroxypropylsulfonates
  - 2-acrylamido-2-methylpropanesulfonic acids
  - metal 2-acrylamido-2-methylpropanesulfonates
  - allylsulfonic acid
  - metal allylsulfonate and the like..

In another embodiment, the polyanionic components include anionic polysaccharides which tend to exist in

ionized forms at higher pH's, for example, pH's of about 7 or higher. The following are some examples of anionic polysaccharides which may be employed in accordance with this invention.

Polydextrose is a randomly bonded condensation polymer of dextrose which is only partially metabolized by mammals. The polymer can contain a minor amount of bound sorbitol, citric acid, and glucose.

Chondroitin sulfate also known as sodium chondroitin sulfate is a mucopolysaccharide found in every part of human tissue, specifically cartilage, bones, tendons, ligaments, and vascular walls. This polysaccharide has been extracted and purified from the cartilage of sharks.

Carrageenan is a linear polysaccharide having repeating galactose units and 3,6 anhydrogalactose units, both of which can be sulfated or nonsulfated, joined by alternating 1-3 and beta 1-4 glycosidic linkages. Carrageenan is a hydrocolloid which is heat extracted from several species of red seaweed and irish moss.

Maltodextrins are water soluble glucose polymers which are formed by the reaction of starch with an acid and/or enzymes in the presence of water.

Other anionic polysaccharides found useful in the present invention are hydrophilic colloidal materials and include the natural gums such as gellan gum, alginate gums, i.e., the ammonium and alkali metal salts of alginic acid and mixtures thereof. In addition, chitosan, which is the common name for deacetylated chitin is useful. Chitin is a natural product comprising poly-(N-acetyl-D-glucosamine). Gellan gum is produced from the fermentation of pseudomonas elodea to yield an extracellular heteropolysaccharide. The alginates and chitosan are available as dry powders from Protan, Inc., Commack, N.Y. Gellan gum is available from the Kelco Division of Merk & Co., Inc., San Diego, Calif.

Generally, the alginates can be any of the water-soluble alginates including the alkali metal alginates, such as sodium, potassium, lithium, rubidium and cesium salts of alginic acid, as well as the ammonium salt, and the soluble alginates of an organic base such as mono-, di-, or triethanolamine alginates, aniline alginates, and the like. Generally, about 0.2% to about 1% by weight and, preferably, about 0.5% to about 3.0% by weight of gellan, alginate or chitosan ionic polysaccharides, based upon the total weight of the composition, are used to obtain the gel compositions of the invention.

Preferably, the anionic polysaccharides are cyclized. More preferably, the cyclized anionic polysaccharides include less than ten monomer units. Even more preferably, the cyclized polysaccharides include less than six monomer units.

In one embodiment, a particularly useful group of cyclized anionic polysaccharides includes the cyclodextrins. Examples of the cyclodextrin group include, but are not limited to: α-cyclodextrin, derivatives of α-cyclodextrin, β-cyclodextrin, derivatives of β-cyclodextrin, γ-cyclodextrin, derivatives of γ-cyclodextrin, carboxymethyl-β-cyclodextrin, carboxymethyl-ethyl-β-cyclodextrin, diethyl-β-cyclodextrin, dimethyl-β-cyclodextrin, methyl-β-cyclodextrin, random methyl-β-cyclodextrin, glucosyl-β-cyclodextrin, maltosyl-β-cyclodextrin, hydroxyethyl-β-cyclodextrin, hydroxypropyl-β-cyclodextrin, sulfobutylether-β-cyclodextrin, and the like and mixtures thereof. Sulfobutylether-β-cyclodextrin is a preferred cyclized anionic polyasaccharide in accordance with the present invention. It is advantageous that the SEC's, including the above mentioned cyclodextrins, employed in this invention be, at the concentration employed, non-toxic to the mammal, human, to inhibit the present incorporation is administered. As used herein, the term "derivatives" as it relates to a cyclodextrin means any substituted or otherwise modified compound which has the characteristic chemical structure of a cyclodextrin sufficiently to function as a cyclodextrin component, for example, to enhance the solubility and/or stability of active components and/or reduce unwanted side effects of the active components and/or to form inclusive complexes with active components, as described herein.

Although cyclodextrins and/or their derivatives may be employed as SECs, one embodimem of the invention may include SECs other than cyclodextrins and/or their derivatives.

A particularly useful and preferred class of polyanionic component includes anionic cellulose derivatives. Anionic cellulose derivatives include metal carboxymethylcelluloses, metal carboxymethylhydroxyethylcelluloses and hydroxypropylmethylcelluloses and derivatives thereof.

The present polyanionic components often can exist in the unionized state, for example, in the solid state, in combination with a companion or counter ion, in particular a plurality of discrete cations equal in number to the number of discrete anionic charges so that the unionized polyanionic component is electrically neutral. For example, the present unionized polyanionic components may be present in the acid form and/or in combination with one or more metals. Since the polyanionic components are preferably ophthalmically acceptable, it is preferred that the metal associated with the unionized polyanionic component be ophthalmically acceptable in the concentrations used. Particularly useful metals include the alkali metals, for example, sodium and potassium, the alkaline earth metals, for example, calcium and magnesium, and mixtures thereof. Sodium is very useful to provide the counter ion in the unionized polyanionic component. Polyanionic components which, in the unionized states, are combined with cations other than H^+ and metal cations can be employed in the present invention.

The amount of SEC in the present compositions is not of critical importance so long as solubility at the alpha-2-adrenergic agonist component is at least somewhat increased and is present in a biologically acceptable amount. Such amount should be effective to perform the desired function or functions in the present composition and/or after administration to the human or animal. In one embodiment, the amount of SEC, preferably the polyanionic component, is sufficient to complex at least in a major amount, and more preferably substantially all, of the alpha-2-adrenergic agonist component in the present composition. In one useful embodiment, the amount of polyanionic component in the present composition is in the range of about 0.1% to about 30% (w/v) or more of the composition. Preferably, the amount of polyanionic component is in the range of about 0.2% (w/v) to about 10% (w/v). More preferably, the amount of polyanionic component is in the range of about 0.2% (w/v) to about 0.6% (w/v). Even more preferably, the polyanionic component is carboxymethylcellulose and is present in the composition in the range of about 0.2% (w/v) to about 0.6% (w/v). A particularly useful concentration of carboxymethylcellulose in the present compositions is about 0.5%.

In one embodiment, the SECs, for example a carboxymethylcellulose, assist in solubilizing the alpha-2-adrenergic agonist components in the compositions. Although the SECs are capable aiding in the solubilization of ionized alpha-2-adrenergic agonist components, it is

preferable that the SECs used in this invention could assist in the solubilization of unionized alpha-2-adrenergic agonist components. For example, in one embodiment, carboxymethylcellulose may help solubilize ionized alpha-2-adrenergic agonist components. In another embodiment, carboxymethylcellulose may help solubilize unionized alpha-2-adrenergic agonist components. In a preferred embodiment, the carboxylmethylcellulose helps solubilize ionized Brimonidine tartrate in the compositions. More preferably, the carboxylmethylcellulose helps solubilize unionized Brimonidine tartrate in the compositions.

In one embodiment, the compositions may also include preservative components or components which assist in the preservation of the composition. The preservative components selected so as to be effective and efficacious as preservatives in the present compositions, that is in the presence of polyanionic components, and preferably have reduced toxicity and more preferably substantially no toxicity when the compositions are administered to a human or animal.

Preservatives or components which assist in the preservation of the composition which are commonly used in pharmaceutical compositions are often less effective when used in the presence of solubilizing agents. In certain instances, this reduced preservative efficacy can be compensated for by using increased amounts of the preservative. However, where sensitive or delicate body tissue is involved, this approach may not be available since the preservative itself may cause some adverse reaction or sensitivity in the human or animal, to whom the composition is administered.

Preferably, the present preservative components or components which assist in the preservation of the composition, preferably the alpha-2-adrenergic agonist components therein, are effective in concentrations of less than about 1% (w/v) or about 0.8% (w/v) and may be 500 ppm (w/v) or less, for example, in the range of about 10 ppm(w/v) or less to about 200 ppm(w/v). Preservative components in accordance with the present invention preferably include, but are not limited to, those which form complexes with the polyanionic component to a lesser extent than does benzalkonium chloride.

Very useful examples of the present preservative components include, but are not limited to oxidative preservative components, for example oxy-chloro components, peroxides, persalts, peracids, and the like, and mixtures thereof. Specific examples of oxy-chloro components useful as preservatives in accordance with the present invention include hypochlorite components, for example hypochlorites; chlorate components, for example chlorates; perchlorate components, for example perchlorates; and chlorite components. Examples of chlorite components include stabilized chlorine dioxide (SCD), metal chlorites, such as alkali metal and alkaline earth metal chlorites, and the like and mixtures therefor. Technical grade (or USP grade) sodium chlorite is a very useful preservative component. The exact chemical composition of many chlorite components, for example, SCD, is not completely understood. The manufacture or production of certain chlorite components is described in McNicholas U.S. Pat. No. 3,278,447, which is incorporated in its entirety herein by reference. Specific examples of useful SCD products include that sold under the trademark Dura Klor by Rio Linda Chemical Company, Inc., and that sold under the trademark Anthium Dioxide by International Dioxide, Inc. An especially useful SCD is a product sold under the trademark Purite™ by Allergan, Inc. Other examples of oxidative preservative components includes peroxy components. For example, trace amounts of peroxy components stabilized with a hydrogen peroxide stabilizer, such as diethylene triamine penta(methylene phosphonic acid) or 1-hydroxyethylidene-1,1-diphosphonic acid, may be utilized as a preservative for use in components designed to be used in the ocular environment. Also, virtually any peroxy component may be used so long as it is hydrolyzed in water to produce hydrogen peroxide. Examples of such sources of hydrogen peroxide, which provide an effective resultant amount of hydrogen peroxide, include sodium perborate decahydrate, sodium peroxide and urea peroxide. It has been found that peracetic acid, an organic peroxy compound, may not be stabilized utilizing the present system. See, for example, Martin et al U.S. Pat. No. 5,725,887, the disclosure of which is incorporated in its entirety herein by reference.

Preservatives other than oxidative preservative components may be included in the compositions. The choice of preservatives may depend on the route of administration. Preservatives suitable for compositions to be administered by one route may possess detrimental properties which preclude their administration by another route. For nasal and ophthalmic compositions, preferred preservatives include quaternary ammonium compounds, in particular the mixture of alkyl benzyl dimethyl ammonium compounds and the like known generically as "benzalkonium chloride." For compositions to be administered by inhalation, however, the preferred preservative is chlorbutol and the like. Other preservatives which may be used, especially for compositions to be administered rectally, include alkyl esters of p-hydroxybenzoic acid and mixtures thereof, such as the mixture of methyl, ethyl, propyl, butyl esters and the like which is sold under the trade name "Nipastat."

In another broad aspect of the present invention, compositions are provided which comprise an alpha-2-adrenergic agonist component, a preservative component in an effective amount to at least aid in preserving, preferably in an amount effective to preserve, the compositions and a liquid carrier component. Preferably, the preservative components include oxy-chloro components, such as compounds, ions, complexes and the like which (1) do not substantially or significantly detrimentally affect the alpha-2-adrenergic agonist components in the compositions or the patients to whom the compositions are administered, and (2) are substantially biologically acceptable and chemically stable. Such compositions in accordance with the present invention comprise an alpha-2-adrenergic agonist component, an oxy-chloro component, and a liquid carrier component, and preferably are substantially free of cyclodextrins.

The carrier components useful in the present invention are selected to be non-toxic and have no substantial detrimental effect on the present compositions, on the use of the compositions or on the human or animal to whom the compositions are administered. In one embodiment, the carrier component is a liquid carrier. In a preferred embodiment, the carrier component is a liquid aqueous carrier component. A particularly useful aqueous liquid carrier component is that derived from saline, for example, a conventional saline solution or a conventional buffered saline solution. The aqueous liquid carrier preferably has a pH in the range of about 6 to about 9 or about 10, more preferably about 6 to about 8, and still more preferably about 7.5. The liquid medium preferably has an ophthalmically acceptable tonicity level, for example, of at least about 200 mOsmol/kg, more preferably in the range of about 200 to about 400 mOsmol/kg. In an especially useful embodiment, the osmolality or tonicity of the carrier component substantially

corresponds to the tonicity of the fluids of the eye, in particular the human eye.

In one embodiment, the carrier components containing the SECs and the alpha-2-adrenergic agonist components may have viscosities of more than about 0.01 centipoise (cps) at 25° C., preferably more than about 1 cps at 25° C., even more preferably more than about 10 cps at 25° C. In a preferred embodiment, the composition has a viscosity of about 50 cps at 25° C. and comprises a conventional buffer saline solution, a carboxymethylcellulose and a Brimonidine tartrate.

In order to insure that the pH of the aqueous liquid carrier component, and thus the pH of the composition, is maintained within the desired range, the aqueous liquid carrier component may include at least one buffer component. Although any suitable buffer component may be employed, it is preferred to select such component so as not to produce a significant amount of chlorine dioxide or evolve significant amounts of gas, such as CO_2. It is preferred that the buffer component be inorganic. Alkali metal and alkaline earth metal buffer components are advantageously used in the present invention.

Any suitable ophthalmically acceptable tonicity component or components may be employed, provided that such component or components are compatible with the other ingredients of the liquid aqueous carrier component and do not have deleterious or toxic properties which could harm the human or animal to whom the present compositions are administered. Examples of useful tonicity components include sodium chloride, potassium chloride, mannitol, dextrose, glycerin, propylene glycol and mixtures thereof. In one embodiment, the tonicity component is selected from inorganic salts and mixtures thereof.

The present compositions may conveniently be presented as solutions or suspensions in aqueous liquids or non-aqueous liquids, or as oil-in-water or water-in-oil liquid emulsions. The present compositions may include one or more additional ingredients such as diluents, flavoring agents, surface active agents, thickeners, lubricants, and the like, for example, such additional ingredients which are conventionally employed in compositions of the same general type.

The present compositions in the form of aqueous suspensions may include excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gun tragacanth and gun acacia; dispersing or wetting agents may be a naturally occurring phosphatide, for example, lecithin, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example, heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol mono-oleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example, polyoxyethylene sorbitan mono-oleate, and the like and mixtures thereof. Such aqueous suspensions may also contain one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose, saccharin, and the like and mixtures thereof.

The present compositions in the form of oily suspensions may be formulated in a vegetable oil, for example, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. Such suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.

The present compositions may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oil, or a mineral oil, for example, liquid paraffin, and the like and mixtures thereof. Suitable emulsifying agents may be naturally-occurring gums, for example, gum acacia or gun tragacanth, naturally-occurring phosphatides, for example, soya bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan mono-oleate, and condensation products of the said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan mono-oleate. The emulsions may also contain sweetening and flavoring agents.

The present compositions in the form of syrups and elixirs may be formulated with sweetening agents, for example, as described elsewhere herein. Such formulations may also contain a demulcent, and flavoring and coloring agents.

The specific dose level for any particular human or animal depends upon a variety of factors including the activity of the active component employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular condition undergoing therapy.

In one broad aspect of the invention, complexes are formed in the present compositions. In one embodiment, the complexes include at least one monomer unit of a quinoxaline component. Examples of quinoxaline components include quinoxaline, (2-imidozolin-2-ylamino) quinoxaline, 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, salts thereof, esters thereof, other derivatives thereof, and the like and mixtures thereof. For example, in one embodiment, a complex of the present invention may include a conjugation of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline monomer units. In another embodiment, the complex may include a conjugation of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline monomer units and Brimonidine tartrate monomer units.

In a preferred embodiment, the complexes of the present invention are dimers. For example, a dimer in accordance with the present invention may include a quinoxaline and a 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline. Preferably, a dimer in accordance with the present invention includes two Brimonidine tartrate monomer units.

Without wishing to limit the invention to any theory or mechanism of operation, it is believed that any peroxide forming agent or strong oxidizing agent such as the oxidative preservative components, for example oxy-chloro components, peroxides, persalts, peracids, and the like, and mixtures thereof may facilitate the formation of the complexes, preferably complexes of alpha-2-adrenergic agonist components. For example, dimers of Brimonidine tartrate monomer units are believed to be formed in the presence of chlorites, preferably stabilized chlorine dioxide.

Furthermore, it is believed that the interactions between the monomers which serve to hold the monomers or monomer subunits together to form a complex, preferably an oligomer and more preferably a dimer, may include, but not limited to, covalent bonding, ionic bonding, hydrophobic bonding, electrostatic bonding, hydrogen bonding, other chemical and/or physical interactions, and the like and combinations thereof. Such complexes may disassociate in liquid, for example, aqueous liquid, media. In one embodiment, the monomers or monomer subunits are held together by other than covalent bonding. In one embodiment, the monomers or monomer subunits are held together by electrostatic bonding or forces.