present invention are hydrophilic colloidal materials and include the natural gums such as gellan gum, alginate gums, i.e., the ammonium and alkali metal salts of alginic acid and mixtures thereof. In addition, chitosan, which is the common name for deacetylated chitin is useful. Chitin is a natural product comprising poly-(N-acetyl-D-glucosamine). Gellan gum is produced from the fermentation of pseudomonas elodea to yield an extracellular heteropolysaccharide. The alginates and chitosan are available as dry powders from Protan, Inc., Commack, N.Y. Gellan gum is available from the Kelco Division of Merk & Co., Inc., San Diego, Calif.

Generally, the alginates can be any of the water-soluble alginates including the alkali metal alginates, such as sodium, potassium, lithium, rubidium and cesium salts of alginic acid, as well as the ammonium salt, and the soluble alginates of an organic base such as mono-, di-, or tri-ethanolamine alginates, aniline alginates, and the like. Generally, about 0.2% to about 1% by weight and, preferably, about 0.5% to about 3.0% by weight of gellan, alginate or chitosan ionic polysaccharides, based upon the total weight of the composition, are used to obtain the gel compositions of the invention.

Preferably, the anionic polysaccharides are cyclized. More preferably, the cyclized anionic polysaccharides include less than ten monomer units. Even more preferably, the cyclized polysaccharides include less than six monomer units.

In one embodiment, a particularly useful group of cyclized anionic polysaccharides includes the cyclodextrins. Examples of the cyclodextrin group include, but are not limited to: α-cyclodextrin, derivatives of α-cyclodextrin, β-cyclodextrin, derivatives of β-cyclodextrin, γ-cyclodextrin, derivatives of γ-

"0071400 0028209

cyclodextrin, carboxymethyl-β-cyclodextrin, carboxymethyl-ethyl-β-cyclodextrin, diethyl-β-cyclodextrin, dimethyl-β-cyclodextrin, methyl-β-cyclodextrin, random methyl-β-cyclodextrin, glucosyl-β-cyclodextrin, maltosyl-β-cyclodextrin, hydroxyethyl-β-cyclodextrin, hydroxypropyl-β-cyclodextrin, sulfobutylether-β-cyclodextrin, and the like and mixtures thereof. Sulfobutylether-β-cyclodextrin is a preferred cyclized anionic polyasaccharide in accordance with the present invention. It is advantageous that the SEC's, including the above mentioned cyclodextrins, employed in this invention be, at the concentration employed, non-toxic to the mammal, human, to inhibit the present incorporation is administered . As used herein, the term "derivatives" as it relates to a cyclodextrin means any substituted or otherwise modified compound which has the characteristic chemical structure of a cyclodextrin sufficiently to function as a cyclodextrin component, for example, to enhance the solubility and/or stability of active components and/or reduce unwanted side effects of the active components and/or to form inclusive complexes with active components, as described herein.

Although cyclodextrins and/or their derivatives may be employed as SECs, one embodiment of the invention may include SECs other than cyclodextrins and/or their derivatives.

A particularly useful and preferred class of polyanionic component includes anionic cellulose derivatives. Anionic cellulose derivatives include metal carboxymethylcelluloses, metal carboxymethylhydroxyethylcelluloses and hydroxypropylmethylcelluloses and derivatives thereof.

The present polyanionic components often can exist in the unionized state, for example, in the solid state, in

CONFIDENTIAL
ALC41213

combination with a companion or counter ion, in particular a plurality of discrete cations equal in number to the number of discrete anionic charges so that the unionized polyanionic component is electrically neutral. For example, the present unionized polyanionic components may be present in the acid form and/or in combination with one or more metals. Since the polyanionic components are preferably ophthalmically acceptable, it is preferred that the metal associated with the unionized polyanionic component be ophthalmically acceptable in the concentrations used. Particularly useful metals include the alkali metals, for example, sodium and potassium, the alkaline earth metals, for example, calcium and magnesium, and mixtures thereof. Sodium is very useful to provide the counter ion in the unionized polyanionic component. Polyanionic components which, in the unionized states, are combined with cations other than H^+ and metal cations can be employed in the present invention.

The amount of SEC in the present compositions is not of critical importance so long as solubility at the alpha-2-adrenergic agonist component is at least somewhat increased and is present in a biologically acceptable amount. Such amount should be effective to perform the desired function or functions in the present composition and/or after administration to the human or animal. In one embodiment, the amount of SEC, preferably the polyanionic component, is sufficient to complex at least in a major amount, and more preferably substantially all, of the alpha-2-adrenergic agonist component in the present composition. In one useful embodiment, the amount of polyanionic component in the present composition is in the range of about 0.1% to about 30% (w/v) or more of the composition. Preferably, the amount of polyanionic component is in the range of about 0.2% (w/v) to about 10%

CONFIDENTIAL
ALC41214

(w/v). More preferably, the amount of polyanionic component is in the range of about 0.2% (w/v) to about 0.6% (w/v). Even more preferably, the polyanionic component is carboxymethylcellulose and is present in the composition in the range of about 0.2% (w/v) to about 0.6% (w/v). A particularly useful concentration of carboxymethylcellulose in the present compositions is about 0.5%.

In one embodiment, the SECs, for example a carboxymethylcellulose, assist in solubilizing the alpha-2-adrenergic agonist components in the compositions. Although the SECs are capable aiding in the solubilization of ionized alpha-2-adrenergic agonist components, it is preferable that the SECs used in this invention could assist in the solubilization of unionized alpha-2-adrenergic agonist components. For example, in one embodiment, carboxymethylcellulose may help solubilize ionized alpha-2-adrenergic agonist components. In another embodiment, carboxymethylcellulose may help solubilize unionized alpha-2-adrenergic agonist components. In a preferred embodiment, the carboxylmethylcellulose helps solubilize ionized Brimonidine tartrate in the compositions. More preferably, the carboxylmethylcellulose helps solubilize unionized Brimonidine tartrate in the compositions.

In one embodiment, the compositions may also include preservative components or components which assist in the preservation of the composition. The preservative components selected so as to be effective and efficacious as preservatives in the present compositions, that is in the presence of polyanionic components, and preferably have reduced toxicity and more preferably substantially no toxicity when the compositions are administered to a human or animal.

CONFIDENTIAL
ALC41215

Preservatives or components which assist in the preservation of the composition which are commonly used in pharmaceutical compositions are often less effective when used in the presence of solubilizing agents. In certain instances, this reduced preservative efficacy can be compensated for by using increased amounts of the preservative. However, where sensitive or delicate body tissue is involved, this approach may not be available since the preservative itself may cause some adverse reaction or sensitivity in the human or animal, to whom the composition is administered.

Preferably, the present preservative components or components which assist in the preservation of the composition, preferably the alpha-2-adrenergic agonist components therein, are effective in concentrations of less than about 1% (w/v) or about 0.8% (w/v) and may be 500 ppm (w/v) or less, for example, in the range of about 10 ppm(w/v) or less to about 200 ppm(w/v). Preservative components in accordance with the present invention preferably include, but are not limited to, those which form complexes with the polyanionic component to a lesser extent than does benzalkonium chloride.

Very useful examples of the present preservative components include, but are not limited to oxidative preservative components, for example oxy-chloro components, peroxides, persalts, peracids, and the like, and mixtures thereof. Specific examples of oxy-chloro components useful as preservatives in accordance with the present invention include hypochlorite components, for example hypochlorites; chlorate components, for example chlorates; perchlorate components, for example perchlorates; and chlorite components. Examples of chlorite components include stabilized chlorine dioxide (SCD), metal chlorites, such as alkali metal and alkaline

0047-0074400

CONFIDENTIAL
ALC41216

earth metal chlorites, and the like and mixtures therefor. Technical grade (or USP grade) sodium chlorite is a very useful preservative component. The exact chemical composition of many chlorite components, for example, SCD, is not completely understood. The manufacture or production of certain chlorite components is described in McNicholas U.S. Patent 3,278,447, which is incorporated in its entirety herein by reference. Specific examples of useful SCD products include that sold under the trademark Dura Klor by Rio Linda Chemical Company, Inc., and that sold under the trademark Anthium Dioxide by International Dioxide, Inc. An especially useful SCD is a product sold under the trademark Purite™ by Allergan, Inc. Other examples of oxidative preservative components includes peroxy components. For example, trace amounts of peroxy components stabilized with a hydrogen peroxide stabilizer, such as diethylene triamine penta(methylene phosphonic acid) or 1-hydroxyethylidene-1,1-diphosphonic acid, may be utilized as a preservative for use in components designed to be used in the ocular environment. Also, virtually any peroxy component may be used so long as it is hydrolyzed in water to produce hydrogen peroxide. Examples of such sources of hydrogen peroxide, which provide an effective resultant amount of hydrogen peroxide, include sodium perborate decahydrate, sodium peroxide and urea peroxide. It has been found that peracetic acid, an organic peroxy compound, may not be stabilized utilizing the present system. See, for example, Martin et al U.S. Patent No. 5,725,887, the disclosure of which is incorporated in its entirety herein by reference.

Preservatives other than oxidative preservative components may be included in the compositions. The choice of preservatives may depend on the route of administration. Preservatives suitable for compositions

CONFIDENTIAL
ALC41217

to be administered by one route may possess detrimental properties which preclude their administration by another route. For nasal and ophthalmic compositions, preferred preservatives include quaternary ammonium compounds, in particular the mixture of alkyl benzyl dimethyl ammonium compounds and the like known generically as "benzalkonium chloride." For compositions to be administered by inhalation, however, the preferred preservative is chlorbutol and the like. Other preservatives which may be used, especially for compositions to be administered rectally, include alkyl esters of p-hydroxybenzoic acid and mixtures thereof, such as the mixture of methyl, ethyl, propyl, butyl esters and the like which is sold under the trade name "Nipastat."

In another broad aspect of the present invention, compositions are provided which comprise an alpha-2-adrenergic agonist component, a preservative component in an effective amount to at least aid in preserving, preferably in an amount effective to preserve, the compositions and a liquid carrier component. Preferably, the preservative components include oxy-chloro components, such as compounds, ions, complexes and the like which (1) do not substantially or significantly detrimentally affect the alpha-2-adrenergic agonist components in the compositions or the patients to whom the compositions are administered, and (2) are substantially biologically acceptable and chemically stable. Such compositions in accordance with the present invention comprise an alpha-2-adrenergic agonist component, an oxy-chloro component, and a liquid carrier component, and preferably are substantially free of cyclodextrins.

The carrier components useful in the present invention are selected to be non-toxic and have no substantial detrimental effect on the present

CONFIDENTIAL
ALC41218

00447070.0028205

compositions, on the use of the compositions or on the human or animal to whom the compositions are administered. In one embodiment, the carrier component is a liquid carrier. In a preferred embodiment, the carrier component is a liquid aqueous carrier component. A particularly useful aqueous liquid carrier component is that derived from saline, for example, a conventional saline solution or a conventional buffered saline solution. The aqueous liquid carrier preferably has a pH in the range of about 6 to about 9 or about 10, more preferably about 6 to about 8, and still more preferably about 7.5. The liquid medium preferably has an ophthalmically acceptable tonicity level, for example, of at least about 200 mOsmol/kg, more preferably in the range of about 200 to about 400 mOsmol/kg. In an especially useful embodiment, the osmolality or tonicity of the carrier component substantially corresponds to the tonicity of the fluids of the eye, in particular the human eye.

In one embodiment, the carrier components containing the SECs and the alpha-2-adrenergic agonist components may have viscosities of more than about 0.01 centipoise (cps) at 25°C, preferably more than about 1 cps at 25°C, even more preferably more than about 10 cps at 25°C. In a preferred embodiment, the composition has a viscosity of about 50 cps at 25°C and comprises a conventional buffer saline solution, a carboxymethylcellulose and a Brimonidine tartrate.

In order to insure that the pH of the aqueous liquid carrier component, and thus the pH of the composition, is maintained within the desired range, the aqueous liquid carrier component may include at least one buffer component. Although any suitable buffer component may be employed, it is preferred to select such component so as not to produce a significant amount of chlorine dioxide or

US021820D.07140D

evolve significant amounts of gas, such as CO_2. It is preferred that the buffer component be inorganic. Alkali metal and alkaline earth metal buffer components are advantageously used in the present invention.

Any suitable ophthalmically acceptable tonicity component or components may be employed, provided that such component or components are compatible with the other ingredients of the liquid aqueous carrier component and do not have deleterious or toxic properties which could harm the human or animal to whom the present compositions are administered. Examples of useful tonicity components include sodium chloride, potassium chloride, mannitol, dextrose, glycerin, propylene glycol and mixtures thereof. In one embodiment, the tonicity component is selected from inorganic salts and mixtures thereof.

The present compositions may conveniently be presented as solutions or suspensions in aqueous liquids or non-aqueous liquids, or as oil-in-water or water-in-oil liquid emulsions. The present compositions may include one or more additional ingredients such as diluents, flavoring agents, surface active agents, thickeners, lubricants, and the like, for example, such additional ingredients which are conventionally employed in compositions of the same general type.

The present compositions in the form of aqueous suspensions may include excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gun tragacanth and gun acacia; dispersing or wetting agents may be a naturally occurring phosphatide, for example, lecithin, or condensation products of ethylene oxide with long chain aliphatic

5021820 - 07400

CONFIDENTIAL
ALC41220

alcohols, for example, heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol mono-oleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example, polyoxyethylene sorbitan mono-oleate, and the like and mixtures thereof. Such aqueous suspensions may also contain one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose, saccharin, and the like and mixtures thereof.

The present compositions in the form of oily suspensions may be formulated in a vegetable oil, for example, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. Such suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.

The present compositions may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oil, or a mineral oil, for example, liquid paraffin, and the like and mixtures thereof. Suitable emulsifying agents may be naturally-occurring gums, for example, gum acacia or gun tragacanth, naturally-occurring phosphatides, for example, soya bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan mono-oleate, and condensation products of the said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan mono-oleate. The emulsions may also contain sweetening and flavoring agents.

The present compositions in the form of syrups and elixirs may be formulated with sweetening agents, for

5025200.07400

CONFIDENTIAL
ALC41221

example, as described elsewhere herein. Such formulations may also contain a demulcent, and flavoring and coloring agents.

The specific dose level for any particular human or animal depends upon a variety of factors including the activity of the active component employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular condition undergoing therapy.

In one broad aspect of the invention, complexes are formed in the present compositions. In one embodiment, the complexes include at least one monomer unit of a quinoxaline component. Examples of quinoxaline components include quinoxaline, (2-imidozolin-2-ylamino) quinoxaline, 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, salts thereof, esters thereof, other derivatives thereof, and the like and mixtures thereof. For example, in one embodiment, a complex of the present invention may include a conjugation of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline monomer units. In another embodiment, the complex may include a conjugation of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline monomer units and Brimonidine tartrate monomer units.

In a preferred embodiment, the complexes of the present invention are dimers. For example, a dimer in accordance with the present invention may include a quinoxaline and a 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline. Preferably, a dimer in accordance with the present invention includes two Brimonidine tartrate monomer units.

Without wishing to limit the invention to any theory or mechanism of operation, it is believed that any peroxide forming agent or strong oxidizing agent such as

CONFIDENTIAL
ALC41222

the oxidative preservative components, for example oxy-chloro components, peroxides, persalts, peracids, and the like, and mixtures thereof may facilitate the formation of the complexes, preferably complexes of alpha-2-adrenergic agonist components. For example, dimers of Brimonidine tartrate monomer units are believed to be formed in the presence of chlorites, preferably stabilized chlorine dioxide.

Furthermore, it is believed that the interactions between the monomers which serve to hold the monomers or monomer subunits together to form a complex, preferably an oligomer and more preferably a dimer, may include, but not limited to, covalent bonding, ionic bonding, hydrophobic bonding, electrostatic bonding, hydrogen bonding, other chemical and/or physical interactions, and the like and combinations thereof. Such complexes may disassociate in liquid, for example, aqueous liquid, media. In one embodiment, the monomers or monomer subunits are held together by other than covalent bonding. In one embodiment, the monomers or monomer subunits are held together by electrostatic bonding or forces.

The following non-limiting examples illustrate certain aspects of the present invention.

5021820.071400

EXAMPLE 1

Brimonidine tartrate has a pKa of about.7.78. The pH-solubility profile of 0.5% (w/v) Brimonidine tartrate in a formulation, Ophthalmic Solution, was established in the pH range of about 5 to about 8 at 23 °C. Table 1. It will be understood that concentrations of adrenergic agonists other than 0.5% may be used, so long as they have therapeutic activity. Likewise, the temperature may be varied, for example, solubility curves may be performed at 37 °C (98.6 °F). The formulation vehicle was prepared by

first dissolving polyvinyl alcohol (PVA) in water. The PVA was added to approximately 1/3 of the required total amount of purified water with constant stirring. The slurry was stirred for 20-30 minutes and then heated to 80-95 °C with constant stirring. The mixture was removed from the heat source within 1 hour after having reached the temperature of 80-90 °C and stirred for an additional 10 minutes to ensure homogeneity (Part I). The other ingredients of the Ophthalmic Solution, except for Brimonidine tartrate, were dissolved in a separate container with an additional 1/3 of the required total amount of purified water (Part II). The PVA mixture (Part I) was then quantitatively transferred to Part II using several rinse volumes of purified water. The solution was adjusted to final volume with purified water without pH adjustment.

Brimonidine tartrate was weighed and transferred to a 10 mL test tube containing 5 mL of the formulation vehicle described above. The pH of each sample was then adjusted to a desired value using dilute sodium hydroxide and/or dilute hydrochloric acid. The samples were placed in a rack on a stir plate and stirred at high speed to achieve uniform mixing for 2 days; a partition was placed between the rack and the stir plate to prevent any heat diffusion from the stir plate to the samples. The temperature of the laboratory was monitored throughout the study and was found to be 23±1 °C.

At the end of two days of stirring, the pH value of each sample was measured, and then approximately 1 mL of each sample was placed in a micro centrifuge tube (polypropylene) and centrifuged at 4,000 rpm for 10 minutes. The supernatant was filtered through a 1μm filter unit (Whatman, 13mm, PTFE). The first 3-4 drops of the filtrate were discarded; the rest of the filtrate was

602182200.07400

received and diluted quantitatively with HPLC mobile phase. The dilute sample was then injected directly on the HPLC column (Dupont Zorbax, 250mm x 4.6mm, 5μm) for Brimonidine tartrate assay in order to quantify the amount of Brimonidine tartrate. A control of 10.05% Brimonidine tartrate was prepared in the formulation vehicle at pH 6.3-6.5 and assayed before (untreated) and after (treated) centrifugation and filtration. This was done to evaluate the potential loss of Brimonidine tartrate in these two steps of the sample preparation. To ensure reproducibility, the study was repeated on consecutive days.

602182000.071400

Table I. 0.5% Brimonidine tartrate in Ophthalmic Solution.

Ingredient	Percent (w/v)
Brimonidine tartrate	0.50
Benzalkonium Chloride, NF	0.0050
Polyvinyl Alcohol, USP	1.4
Sodium Chloride, USP	0.66
Sodium Citrate, Dihydrate, USP	0.45
Hydrochloric Acid, NF or Sodium Hydroxide, NF for pH adjustment	5-8
Purified Water, USP	QS

The solubility data for Brimonidine tartrate in the formulation vehicles are presented in Table II. The results show that the solubility of Brimonidine tartrate is highly pH-dependent and spans more than two orders of magnitude over the pH range of 5-8. The solubility decreases sharply as the pH increases. The results for the treated and untreated controls are very close, suggesting that centrifugation and filtration does not cause any significant loss of Brimonidine tartrate. The

CONFIDENTIAL
ALC41225

two solubility profiles obtained on consecutive days agree with each other.

0021202.0714.08

CONFIDENTIAL
ALC41226

Table II. Solubility of Brimonidine tartrate in the Ophthalmic Solution Over pH Range of 5 to 8.

	STUDY 1		STUDY 2	
Sample	pH[a]	Solubility[a]	pH[a]	Solubility[a]
1	5.55	≥164.4[b]	5.50	≥200.6[b]
2	5.92	132.6	5.92	160.8
3	6.14	30.4	6.06	50.1
4	6.57	7.55	6.90	3.19
5	7.00	2.69	7.40	1.19
6	7.45	1.17	7.77	0.63
7	7.83	0.62	7.86	0.58
8	—	—	7.88	0.54
Control/ (untreated)	—	0.486[c]	—	—
Control/ (treated)	—	0.484[d]	—	—

[a] Measured after stirring for two-days before sample withdrawal for centrifugation and filtration.

[b] Represents theoretical concentration based on sample weight. The sample solution was clear indicating that all of the Brimonidine tartrate had dissolved.

[c] Concentration of Brimonidine tartrate in control before centrifugation and filtration step.

[d] Concentration of Brimonidine tartrate in control after centrifugation and filtration step.

[e] %W/V.

00147.0002.5028209

EXAMPLE 2

The pH-solubility profiles of Brimonidine tartrate in compositions (solutions) containing SECs and oxy-chloro components were determined. Particularly, the effects of sodium carboxymethylcellulose (CMC), an SEC, on the solubility of Brimonidine tartrate at various pH conditions were determined. The various concentrations of CMC tested with Brimonidine tartrate were 0%, 0.056%, 0.17%, 0.5%, 1.5% (w/v), Table III.

The samples tested also contained isotonic components, buffer components, and stabilized chlorine dioxide (Purite™), Table III. Sodium carboxymethylcellulose, sodium chloride, potassium chloride, calcium chloride dihydrate, and magnesium chloride hexahydrate were USP grade. Boric acid and sodium borate decahydrate were NF grade.

Table III

	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	
Brimonidine tartrate	0.2%	0.2%	0.2%	0.2%	0.2%	(w/v)
CMC	0.0%	0.056%	0.17%	0.5%	1.5%	(w/v)
Stabilized chlorine dioxide[a]	0.005%	0.005%	0.005%	0.005%	0.005%	(w/v)
Sodium chloride	0.58%	0.58%	0.58%	0.58%	0.58%	(w/v)
Potassium chloride	0.14%	0.14%	0.14%	0.14%	0.14%	(w/v)
Calcium chloride, dihydrate	0.02%	0.02%	0.02%	0.02%	0.02%	(w/v)
Magnesium chloride, hexahydrate	0.006%	0.006%	0.006%	0.006%	0.006%	(w/v)
boric acid	0.2%	0.2%	0.2%	0.2%	0.2%	(w/v)
sodium tetraborate, decahydrate	0.14%	0.14%	0.14%	0.14%	0.14%	(w/v)

[a] Sold under the trademark Purite™ by Allergan, Inc.

CONFIDENTIAL
ALC41228

Each sample (1 through 5) was subjected to a range of pH's from about 7 to about 10. The vials containing the sample solutions were placed on a laboratory rotator and left for equilibration for fifteen days at room temperature (~21 °C). The sample solutions were filtered using a 25 mm diameter polysulfone cellulose acetate syringe type filter with 0.45µm pore size. The filtered solutions were assayed for Brimonidine.

Conventional HPLC and detection techniques were used to detect and determine the concentrations of soluble Brimonidine tartrate. Table IV. The solubility is plotted against pH for each CMC concentration. The experimental data points were fitted to a modified Henderson-Hasselbalch equation using a nonlinear least squares routine (Deltagraph version 4.0 DeltaPoint, Inc.), Fig. 1. The R^2 values show the goodness of fit between the experimental values and the theoretical equation to be better than 0.991.

0021800.0714.00

CONFIDENTIAL
ALC41229

Table IV

		Solubility of Brimonidine tartrate (%)				
		0%CMC	0.056%CMC	0.17%CMC	0.5%CMC	1.5%CMC
	pH					
5	6.67		0.9302		1.4464	
	6.68	1.4256		1.4200		
	6.93			0.7302		
	7.10				0.3693	
	7.11	0.2064	0.2828			
10	7.35					0.1904
	7.56				0.1451	
	7.68	0.0786				
	7.77		0.0721			
	7.81			0.0735		
15	8.10					0.0498
	8.46				0.0313	
	8.50	0.0286				
	8.55			0.0328		
	8.67					0.0311
20	9.93		0.0234			
	9.94				0.0250	
	10.05			0.0241		
	10.09	0.0218				
	10.11					0.0222

CONFIDENTIAL
ALC41230

Fig. 1 clearly shows that the solubility of Brimonidine tartrate tends to increase with increasing CMC concentrations. For example, at pH 7.5, the sample with 0% CMC resulted in 1000 ppm of Brimonidine tartrate; 0.056% CMC, 1300 ppm; 0.17% CMC, 1300 ppm; and 0.5%, 1600 ppm. At pH 7.5, the sample with 1.5% CMC resulted in about 1400 ppm, which is less than that of a similar solution with CMC at 0.5%. It is unclear at this point what the cause of this observation may be. Nonetheless, Brimonidine tartrate is more soluble in solution with a 1.5% CMC than with no CMC.

CMC is also effective to solubilize Brimonidine tartrate in a biological environment, for example the biological environment of the cornea.

EXAMPLE 3

Brimonidine tartrate dimers.

Brimonidine tartrate is added to a test tube containing a composition including chlorite. The test tube was allowed to equilibrate for ten days. Samples obtained from the test tube is analyzed. It is observed that a portion of the Brimonidine tartrate monomer units conjugated to form dimers.

While this invention has been described with respect to various specific examples and embodiments, it is to be understood that the invention is not limited thereto and that it can be variously practiced with the scope of the following claims.

0021820.0714.00

CONFIDENTIAL
ALC41231

WHAT IS CLAIMED IS:

1. A composition comprising:

an alpha-2-adrenergic agonist component in an amount effective to provide a therapeutic benefit to a patient to whom the composition is administered;

a solubility enhancing component in an amount effective to increase the solubility of the alpha-2-adrenergic agonist component in the composition relative to the solubility of an identical alpha-2-adrenergic agonist component in a similar composition without the solubility enhancing component; and

a liquid carrier component.

2. The composition of claim 1 wherein the alpha-2-adrenergic agonist component is selected from the group consisting of imino-imidazolines, imidazolines, imidazoles, azepines, thiazines, oxazolines, guanidines, catecholamines, derivatives thereof and mixtures thereof.

3. The composition of claim 1 wherein the therapeutically active component includes a quinoxaline component.

4. The composition of claim 3 wherein the quinoxaline component is selected from the group consisting of quinoxaline, derivatives thereof, and mixtures thereof.

5. The composition of claim 3 wherein the quinoxaline component is selected from the group consisting of quinoxaline, (2-imidozolin-2-ylamino) quinoxaline, 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, and tartrate of 5-bromo-6-(2-imidozolin-2-

CONFIDENTIAL
ALC41232

ylamino) quinoxaline, derivatives thereof and mixtures thereof.

6. The composition of claim 1 wherein the therapeutically active component comprises a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline.

7. The composition of claim 1 wherein the alpha-2-adrenergic agonist component is substantially unionized.

8. The composition of claim 1 wherein the alpha-2-adrenergic agonist component is substantially unionized in a biological environment to which the composition is administered.

9. The composition of claim 1 wherein the alpha-2-adrenergic agonist component has increased diffusion through a lipid membrane relative to an identical alpha-2-adrenergic agonist component in a similar composition without the solubility enhancing component.

10. The composition of claim 1 wherein the alpha-2-adrenergic agonist component is selected from the group consisting of agonists of alpha-2A-adrenergic receptors, agonists of alpha-2B-adrenergic receptors, agonists of alpha-2D-adrenergic receptors and mixtures thereof.

11. The composition of claim 1 wherein the solubility enhancing component is effective to increase the solubility in a biological environment of the alpha-2-adrenergic agonist component relative to the solubility in a biological environment of an identical alpha-2-adrenergic agonist component in a similar composition without the solubility enhancing component.

CONFIDENTIAL
ALC41233

6021200.07400

12. The composition of claim 1 wherein the solubility enhancing component comprises a polyanionic component.

13. The composition of claim 12 wherein said polyanionic component is selected from the group consisting of anionic cellulose derivatives, anionic polymers derived from acrylic acid, anionic polymers derived from methacrylic acid, anionic polymers derived from alginic acid, anionic polymers derived from amino acids and mixtures thereof.

14. The composition of claim 1 wherein the solubility enhancing component is selected from the group consisting of anionic cellulose derivatives and mixtures thereof.

15. The composition of claim 1 wherein the solubility enhancing component is selected from the group consisting of carboxymethylcelluloses and derivatives thereof.

16. The composition of claim 1 wherein the solubility enhancing component is present in an amount in a range of about 0.1% (w/v) to about 30% (w/v).

17. The composition of claim 1 wherein the solubility enhancing component is present in an amount in a range of about 0.2% (w/v) to about 10% (w/v).

18. The composition of claim 1 wherein the solubility enhancing component is present in an amount in a range of about 0.2% (w/v) to about 0.6% (w/v).

0044.0 07128200

CONFIDENTIAL
ALC41234

19. The composition of claim 1 wherein the liquid carrier component is an aqueous liquid carrier component.

20. The composition of claim 1 which is a solution.

21. The composition of claim 1 which has a pH of about 7 or greater.

22. The composition of claim 1 which has a pH in a range of about 7 to about 9.

23. The composition of claim 1 which is ophthalmically acceptable.

24. A composition comprising:

an alpha-2-adrenergic agonist component in an amount effective to provide a therapeutic benefit to a patient to whom the composition is administered;

an anionic cellulose derivative in an amount effective to increase the solubility of the alpha-2-adrenergic agonist component; and

an aqueous liquid carrier component.

25. The composition of claim 24 wherein the alpha-2-adrenergic agonist component comprises a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline.

26. The composition of claim 24 wherein the anionic cellulose derivative comprises carboxymethylcellulose.

27. The composition of claim 24 wherein the anionic cellulose derivative is present in an amount in a range of about 0.2% (w/v) to about 0.6% (w/v).

5028220.0704

CONFIDENTIAL
ALC41235

28. A composition comprising:

a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline in an amount effective to provide a therapeutic benefit to a patient to whom the composition is administered;

a solubility enhancing component in an amount effective to increase the solubility of the tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline; and

an aqueous liquid carrier component.

29. The composition of claim 28 wherein the solubility enhancing component comprises a carboxymethylcellulose.

30. The composition of claim 28 which is ophthalmically acceptable.

31. A complex comprising monomer units derived from one or more quinoxaline components.

32. The complex of claim 31 wherein the quinoxaline component is selected from the group consisting of a quinoxaline, a (2-imidozolin-2-ylamino) quinoxaline, a 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, derivatives thereof and mixtures thereof.

33. The complex of claim 31 wherein the quinoxaline component is a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline.

34. An oligomer comprising monomer units derived from a quinoxaline component.

CONFIDENTIAL
ALC41236

ALC41236 - 00281209

35. The oligomer of claim 34 wherein the quinoxaline component is selected from the group consisting of a quinoxaline, a (2-imidozolin-2-ylamino) quinoxaline, a 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline, derivatives thereof and mixtures thereof.

36. The oligomer of claim 34 wherein the quinoxaline component is a tartrate of 5-bromo-6-(2-imidozolin-2-ylamino) quinoxaline.

37. The oligomer of claim 34 which is a dimer.

38. The oligomer of claim 35 which is a dimer.

39. The oligomer of claim 36 which is a dimer.

40. A composition comprising:

an alpha-2-adrenergic agonist component in an amount effective to provide a therapeutic benefit to a patient to whom the composition is administered;

an oxy-chloro component in an effective amount to at least aid in preserving the composition; and

a liquid carrier component,

wherein the composition is substantially free of cyclodextrins.

41. The composition of claim 40 wherein the alpha-2-adrenergic agonist component is selected from the group consisting of imino-imidazolines, imidazolines, imidazoles, azepines, thiazines, oxazolines, guanidines, catecholamines, derivatives thereof and mixtures thereof.

42. The composition of claim 40 wherein the

CONFIDENTIAL
ALC41237

therapeutically active component includes a quinoxaline component.

43. The composition of claim 42 wherein the quinoxaline component is selected from the group consisting of quinoxaline, derivatives thereof, and mixtures thereof.


44. The composition of claim 40 which further includes a solubility enhancing component in an amount effective to increase the solubility of the alpha-2-adrenergic agonist component in the composition relative to the solubility of an identical alpha-2-adrenergic agonist component in a similar composition without the solubility enhancing component.

45. The composition of claim 44 wherein the solubility enhancing component is effective to increase the solubility in a biological environment of the alpha-2-adrenergic agonist component relative to the solubility in a biological environment of an identical alpha-2-adrenergic agonist component in a similar composition without the solubility enhancing component.


46. The composition of claim 44 wherein the solubility enhancing component comprises a polyanionic component.

5021820.07400

Figure 1



CONFIDENTIAL
ALC41239



CONFIDENTIAL
ALC41240

PTO/SB/58 (07-03)
Approved for use through 7/31/2003. OMB 0651-0031
U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

REQUEST FOR ACCESS TO AN ABANDONED APPLICATION UNDER 37 CFR 1.14

Bring completed form to:
File Information Unit
Crystal Plaza Three, Room 1D01
2021 South Clark Place
Arlington, VA
Telephone: (703) 308-2733

RECEIVED
DEC 0 3 2003
File Information Unit

In re Application of

Application Number	Filed
60/218200	7-12-00

Paper No. #2

I hereby request access under 37 CFR 1.14(a)(1)(iv) to the application file record of the above-identified ABANDONED application, which is identified in, or to which a benefit is claimed, in the following document (as shown in the attachment):

United States Patent Application Publication No. _____, page, _____ line _____,

United States Patent Number 6627210, column _____, line, _____ or

WIPO Pub. No._____, page _____, line _____.

Related Information about Access to Pending Applications (37 CFR 1.14):
Direct access to pending applications is not available to the public but copies may be available and may be purchased from the Office of Public Records upon payment of the appropriate fee (37 CFR 1.19(b)), as follows:
For published applications that are still pending, a member of the public may obtain a copy of:
the file contents;
the pending application as originally filed; or
any document in the file of the pending application.
For unpublished applications that are still pending:
(1) If the benefit of the pending application is claimed under 35 U.S.C. 119(e), 120, 121, or 365 in another application that has: (a) issued as a U.S. patent, or (b) published as a statutory invention registration, a U.S. patent application publication, or an international patent application publication in accordance with PCT Article 21(2), a member of the public may obtain a copy of:
the file contents;
the pending application as originally filed; or
any document in the file of the pending application.
(2) If the application is incorporated by reference or otherwise identified in a U.S. patent, a statutory invention registration, a U.S. patent application publication, or an international patent application publication in accordance with PCT Article 21(2), a member of the public may obtain a copy of:
the pending application as originally filed.

Signature

12-3-03
Date

Typed or printed name

Registration Number, if applicable

(703) 415-1671
Telephone Number

FOR PTO USE ONLY
Approved by:
Unit:
RECEIVED
DEC 0 3 2003
File Information Unit

This collection of information is required by 37 CFR 1.14. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS

CONFIDENTIAL
ALC41241

12/02/03 TUE 13:35 FAX 18006661233 001

US006627210B2

(12) **United States Patent**
Olejnik et al.

(10) Patent No.: US 6,627,210 B2
(45) Date of Patent: *Sep. 30, 2003

(54) **COMPOSITIONS CONTAINING α-2-ADRENERGIC AGONIST COMPONENTS**

(75) Inventors: Orest Olejnik, Coto de Caza, CA (US); Edward D. S. Karslaka, Charlestown, MA (US)

(73) Assignee: Allergan, Inc., Irvine, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 8 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: 09/904,918

(22) Filed: Jul. 10, 2001

(65) Prior Publication Data

US 2002/0022201 A1 Mar. 14, 2002

Related U.S. Application Data

Provisional application No. 60/218,200, filed on Jul. 14, 2000.

(51) Int. Cl.[7] A61K 33/14; A61K 9/00; A61K 9/08

(52) U.S. Cl. 424/427; 424/400; 424/401; 424/466; 424/422; 514/772.4; 514/772.6

(58) Field of Search 424/427, 400, 424/422, 661, 407; 514/772.4, 772.6

(56) References Cited

U.S. PATENT DOCUMENTS

3,270,447	A	10/1966	McGoldrick	
3,890,319	A	6/1975	Czarkowski et al.	
4,530,920	A	7/1985	Nesler et al.	
4,806,556	A	2/1989	Portoghese	
5,021,416	A	6/1991	Czarkowski	
5,202,128	A	4/1993	Morella et al.	
5,215,991	A *	6/1993	Burke	514/255
5,352,796	A	10/1994	Hooper et al.	
5,459,130	A *	10/1995	Neufeld	514/211
5,708,077	A	12/1997	Burke et al.	
5,718,197	A	2/1998	Keates et al.	
5,725,887	A	3/1998	Martin et al.	
5,814,638	A	9/1998	Lee et al.	
5,834,502	A	11/1998	Cleary et al.	
5,994,110	A	11/1999	Meebach et al.	
2002/0022202	A1 *	1/2002	Gant	514/892
2002/0071874	A1 *	6/2002	Olejnik et al.	424/451

FOREIGN PATENT DOCUMENTS

CA	2048315	2/1992
EP	0509961	8/1994
WO	94/15585	8/1994
WO	98/47878	10/1998
WO	99/43299	9/1999
WO	99/51273	10/1999
WO	00/12157	3/2000
WO	00/19981	4/2000

OTHER PUBLICATIONS

Remington's Pharmaceutical Sciences, Eighteenth Edition, 1990, pp. 1504–1505.*
U.S. patent application Ser. No. 09/903,962, Olejnik et al., filed Jul. 10, 2001.


* cited by examiner

Primary Examiner—Thurman K. Page
Assistant Examiner—Rachel M. Bennett
(74) *Attorney, Agent, or Firm*—Carlos A. Fisher; Martin A. Voss; Robert J. Baran

(57) ABSTRACT

Compositions useful for improving effectiveness of alpha-2-adrenergic agonist components include carrier components, alpha-2-adrenergic agonist components, solubility enhancing components which aid in solubilizing the alpha-2-adrenergic agonist components. In one embodiment, the alpha-2-adrenergic agonist components include alpha-2-adrenergic agonists. In another embodiment, the solubility enhancing components include carboxymethylcelluloses.

34 Claims, 1 Drawing Sheet



CONFIDENTIAL
ALC41242

CERTIFICATE OF SERVICE

I, Karen E. Keller, Esquire hereby certify that on May 2, 2005, I caused to be electronically filed a true and correct copy of the foregoing document with the Clerk of the Court using CM/ECF, which will send notification that such filing is available for viewing and downloading to the following counsel of record:


William J. Marsden, Jr., Esquire
Fish & Richardson, P.A.
919 N. Market Street, Suite 1100
Wilmington, DE 19801

I further certify that on May 2, 2005, I caused a copy of the foregoing document to be served by hand delivery on the above-listed counsel of record and on the following non-registered participants in the manner indicated:

BY FEDERAL EXPRESS

Jonathan E. Singer, Esquire
Fish & Richardson, P.A.
3300 Dain Rauscher Plaza
60 South Sixth Street
Minneapolis, MN 55402

YOUNG CONAWAY STARGATT & TAYLOR, LLP



Josy W. Ingersoll (No. 1088)
Karen E. Keller (No. 4489)
The Brandywine Building
1000 West Street, 17th Floor
Wilmington, Delaware 19801
(302) 571-6600
kkeller@ycst.com

Attorneys for Defendants